



Received on 10 September, 2018; received in revised form, 24 September, 2018; accepted, 22 October, 2018; published 01 November, 2018

DIETARY POLYPHENOLS: AN OVERVIEW

A. P. Attanayake

Department of Biochemistry, Faculty of Medicine, University of Ruhuna, Galle, Sri Lanka.

ABSTRACT: Dietary polyphenols, mainly phenolic acids and flavonoids, are the primary sources of antioxidants that can be generally recommended for human consumption. Polyphenols are derived from plant based food including fruits, vegetables, spices, herbs *etc.* In the present era, there has been a great interest towards the health benefits of dietary polyphenols against oxidative stress related diseases such as diabetes mellitus, cancer, chronic kidney disease *etc.* Here, the content presents an overview on dietary polyphenols in context of relevance to the human health.

Keywords: Antioxidant activity, Chronic diseases, Polyphenols

Correspondence to Author:

Dr. A. P. Attanayake

Department of Biochemistry, Faculty of Medicine, University of Ruhuna, Galle, Sri Lanka.

E-mail: anoja715@yahoo.com

INTRODUCTION: Polyphenols are the largest group of phytonutrients that exert different biological activities that are beneficial to human health. Fruits, vegetables, cereals, and plant-derived beverages such as fruit juices, tea, coffee and red wine are the main dietary sources of polyphenols. It is found that human consumption of polyphenols is, generally higher than any other phytonutrient or dietary antioxidant^{1, 2, 3}. The scientific and research interest in biological activities of polyphenols for maintaining human health approximately started in late 1980s and a tremendous progress has been made to date^{4, 5}. Polyphenols, are widely distributed in the plant kingdom. More than 8,000 distinct polyphenol structures are reported and identified from different plants⁶.

Polyphenols are the products of secondary metabolism and can vary from simple phenolic acids to highly polymerized molecules such as tannins. Polyphenols exist primarily in conjugated forms with one or more sugar residues, via glycosides, linked to one or more of the many hydroxyl groups^{4, 7}. Polyphenols can be classified into groups according to their source of origin such as tea polyphenols, grape polyphenols and apple polyphenols, or their chemical structures. When basic chemical structure is of concern, polyphenols can be divided into basic groups as lignans, phenolic acids, flavonoids, stilbenes⁸.

The most common and important low molecular weight phenolic compounds are simple phenolic derivatives and flavonoids⁹. Phenolic acids are simple molecules such as caffeic acid, vanillin and coumaric acid that account for approximately one-third of the total dietary intake of polyphenols. Many flavonoids present in plants as aglycones although they are most commonly found as glycoside derivatives. Flavonoids account for the remaining two-thirds. Flavonoids can be further subdivided into classes including flavonols,

	QUICK RESPONSE CODE DOI: 10.13040/IJPSR.0975-8232.IJLSR.4(11).176-79
	Article can be accessed online on: www.ijlsr.com
DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.IJLSR.4(11).176-79	

flavanones, flavanols, flavones, anthocyanins, isoflavones with more than 4,000 described compounds^{10, 11, 12}.

Occurrence: Polyphenol compounds are distributed in plants at the tissue, cellular and sub cellular levels. However, the distribution of polyphenols is not uniform. Insoluble polyphenols are mainly found in cell walls; however soluble polyphenols are present within the plant cell vacuoles¹³. Numerous factors affect the polyphenol content of plants, such as climatic and environmental factors, processing, storage, degree of ripeness at the time of harvest, *etc.* The outer layers of plants contain higher levels of polyphenols than those located in the inner parts¹⁰.

In most cases, food contains complex mixtures of polyphenols¹⁴. Flavonoids are mostly found in all plant based food; fruits, vegetables, cereals, fruit juices, tea, wine, infusions *etc.*, whereas flavanones and isoflavones are specific to particular foods¹⁵. The degree of ripeness considerably affects the concentrations and proportions of various polyphenols^{16, 17}. The easy oxidation of the polyphenols may lead to change the content of polyphenols in plants¹⁶. Different types of cooking methods also affect the polyphenol content of foods since polyphenols are present in plant vacuoles and in cellular structures that require softening during the cooking process to maximize polyphenol release. However, most of the vegetables can lose 75% to 100% of their initial polyphenol content during boiling whereas steaming and frying losses are considerably lower than fresh vegetables¹⁸.

Bioavailability: Dietary intake of plant polyphenols can vary considerably and be quite high. However, ultimately the nutritional impact and subsequent systemic effects depend on the fate of polyphenols in the human digestive tract^{19, 20}. Physicochemical characteristics of polyphenols appear to dictate absorption and metabolism and are determined primarily by the chemical structures^{6, 7}. Relatively small molecular weight phenolic acids, *i.e.*, gallic acid and isoflavones are easily absorbed through the gut followed by catechins, flavanones, and quercetin glucosides. Larger polyphenols such as proanthocyanidins are very poorly absorbed. The main difficulty with incorporating polyphenols into existing products is

to present them in a form that is bioavailable²¹. Accordingly, the total polyphenol metabolite content in plasma or in urine after intake of dietary polyphenols is generally low^{2, 22}. It was further noted that bioavailability is differed greatly from one polyphenol to another and the most abundant dietary polyphenol was not necessarily the one leading to the highest levels of active metabolites in plasma. The maximum plasma concentrations of polyphenol metabolite content is less than 10 $\mu\text{mol/L}$ ²³. The physicochemical characteristics of polyphenols likely contribute to the disparity in results between the epidemiological studies and *in-vitro* data, which may reflect, in part, relatively poor bioavailability, rapid metabolism, and excretion of polyphenols²⁴.

Scientists are being taking efforts to increase the bioavailability of polyphenols but several considerations, and potential impediments, exist including solubility, permeability, metabolism, excretion, target tissue uptake using nanoparticle based approaches and disposition²¹. Initial strategies for improving bioavailability of dietary polyphenol supplements include changes to polyphenol structure, which may affect the solubility and dissolution, and the use of pharmaceutical recipients²¹. An additional consideration is that degradation and absorption of polyphenols within the gastrointestinal tract depend on the intestinal microflora and gut enzymes, which may significantly change bioavailability²⁵.

Metabolism of Polyphenols: Dietary polyphenols are initially metabolized, within the gastrointestinal tract. Glycosides are hydrolyzed to their corresponding aglycones prior to the absorption, aglycones and low molecular weight polyphenols are absorbed directly². Polyphenols that are not absorbed are transported to the colon where they can be readily hydrolyzed by colonic microflora to simpler chemical compounds and eliminated in the feces or further modified. Bacterial fermentation of the glycones effectively releases the polyphenol from fiber permitting subsequent metabolism similar to other aglycones²⁶.

Polyphenols and Human Diseases: Correlations between the intake of polyphenols and reduced incidence of chronic diseases have been firmly established using different *in-vivo* models in the

past decades. The in cooperation of polyphenols in the prevention and management of inflammation and oxidative stress related pathologies is emerging⁴. Epidemiological studies have repeatedly shown an inverse association between the occurrence of chronic diseases and the consumption of polyphenolic rich diet^{27, 28}. It is well established that polyphenol-rich foods and beverages may increase plasma antioxidant capacity². This increase in the antioxidative capacity of plasma following the consumption of polyphenol-rich food may be explained either by the presence of reducing polyphenols and their metabolites in plasma, by their effects upon concentrations of other reducing agents *via* sparing effects of polyphenols on other endogenous antioxidants, or by their effect on the absorption of pro-oxidative food components, such as iron²⁹. Consumption of antioxidants has been associated with reduced levels of oxidative damage to lymphocytic DNA³⁰. There are increasing evidences that as antioxidants, polyphenols may protect cell constituents against oxidative damage and therefore, limit the risk of various degenerative diseases associated with oxidative stress^{31, 32}.

CONCLUSION: The content outlined in this review provides a general overview of dietary polyphenols, occurrence and content, bio-availability, metabolism and their relevance to human health. Dietary polyphenols are beneficial against the development and progression of many chronic oxidative stress related pathological conditions. The role of polyphenols in human health and disease is still an interesting area of research and warranted further scientific studies.

ACKNOWLEDGEMENT: Nil

CONFLICT OF INTEREST: Nil

REFERENCES:

1. Scalbert AG and Williamson: Dietary intake and bioavailability of polyphenols. *Journal of Nutrition* 2000; 130: 2073S-2078S.
2. Scalbert A, Johnson IT and Saltmarsh M: Polyphenols: antioxidants and beyond. *American Journal of Clinical Nutrition* 2005; 81: 215S-217S.
3. Williamson G: The role of polyphenols in modern nutrition. *Nutrition Bulletin* 2017; 42(3): 226-235.
4. Shen T, Han X, Wang X, Fan P, Ren D and Lou H: Protective effects of dietary polyphenols in human diseases and mechanisms of action. In *Nutritional antioxidant therapies: Treatments and Perspectives*. Springer International Publishing AG. Switzerland 2007; 307-346.
5. Adebooye OC, Alashi AM and Aluko RE: A brief review on emerging trends in global polyphenol research. *Journal of Food Biochemistry* 2018; e12519.
6. Rio DD, Rodriguez-Mateos A, Spencer JPE, Borges T and Crozier A: Dietary polyphenolics in human health: Structures, bioavailability and evidence of protective effects against chronic diseases. *Antioxidants and Redox Signaling* 2013; 18(14): 1818-1892.
7. Martin KR and Appel CL: Polyphenols as dietary supplements: A double-edged sword. *Nutrition and Dietary Supplements* 2010; 2: 1-12.
8. Crozier A, Jaganath IB and Clifford MN: Dietary phenolics: Chemistry, new magic bullet? *Proceedings of the Nutrition Society* 2003; 62: 599-603.
9. Alkhatib A, Tsang C, Tiss A, Bahorun T, Arefanian H, Barake R, Khadir A and Tuomilehto J: Functional foods and lifestyle approaches for diabetes prevention and management. *Nutrients* 2017; 9: 1310.
10. Pandey KB and Rizvi SI: Plant polyphenols as dietary antioxidants in human health and disease. *Oxidative Medicine and Cell Longevity* 2009; 2(5): 270-278.
11. Kumar S and Pandey AK: Chemistry and biological activities of flavonoids: An overview. *Scientific World Journal* 2013; e162750.
12. Carmona-Hernandez JC, Ceballos-López LM and González-Correa CH: Polyphenols and flavonoids in Colombian fruit and vegetables - applications and benefits: A review. *Journal of Food Nutrition Research* 2018; 6 (3): 76-181.
13. Wink M: Compartmentation of secondary metabolites and xenobiotics in plant vacuoles. *Advances in Botanical Research* 1997; 25: 141-169.
14. Alkhalidy HY and Liu WD: Dietary flavonoids in the prevention of T2D: An overview. *Nutrients* 2018; 10: 438.
15. Gharras HEI: Polyphenols: food sources, properties and applications - A review. *International Journal of Food Science and Technology* 2009; 44: 251-2518.
16. Manach C, Scalbert A, Morand C, Rémésy C and Jimenez L: Polyphenols: food sources and bioavailability. *American Journal of Clinical Nutrition* 2004; 79: 727-747.
17. Simon BF, Perez-Illzarbe J, Hernandez T, Gomez-Cordoves C and Estrella I: Importance of phenolic compounds for the characterization of fruit juices. *Journal of Agricultural and Food Chemistry* 1992; 40: 1531-1535.
18. Miglio C, Chiavaro E, Visconti A, Fogliano V and Pellegrini N: Effects of different cooking methods on nutritional and physicochemical characteristics of selected vegetables. *Journal of Agricultural and Food Chemistry* 2008; 56: 139-147.
19. Ovaskainen M, Torronen R, Koponen JM, Sinkko H, Hellstrom J, Reinivuo H and Mattila P: Dietary intake and major food sources of polyphenols in Finnish adults. *Journal of Nutrition* 2008; 138: (3): 562-566.
20. Santhakumara AB, Battinob M and Alvarez-Suarezc MJM: Dietary polyphenols: Structures, bioavailability and protective effects against atherosclerosis. *Food and Chemical Toxicology* 2018; 113: 49-65.
21. Hu M: Commentary: Bioavailability of flavonoids and polyphenols: Call to arms *Molecular Pharmaceutics* 2007; 4: 803-806.
22. Jew S, Suhad S, Mweis A and Jones PJH: Evolution of the human diet: Linking our ancestral diet to modern functional foods as a means of chronic disease prevention. *Journal of Medicinal Food* 2009; 12 (5): 925-934.

23. Kroon PA, Clifford MN, Crozier A, Day AJ, Donovan JL, Manach C and Williamson G: How should we assess the effects of exposure to dietary polyphenols *in-vitro*? American Journal of Clinical Nutrition 2004; 80(1): 15-21.
24. Duthie GG, Gardner PT and Kyle JAM: Plant polyphenols: are they the new magic bullet? 2003; 62(3): 599-603.
25. Marín L, Miguélez EM, Villar CJ and Lombó F: Bioavailability of dietary polyphenols and gut microbiota metabolism: Antimicrobial properties. Biomed Research International 2015; ID 905215.
26. Bravo L: Polyphenols: Chemistry, dietary sources, metabolism, and nutritional significance. Nutrition Reviews 1998; 56: 317-333.
27. Arts ICW and Hollman PC: Polyphenols and disease risk in epidemiologic studies. American Journal of Clinical Nutrition 2005; 81: 317-325.
28. Clifford MN: Chlorogenic acids and other cinnamates. Nature, occurrence, dietary burden, absorption and metabolism. Journal of Science of Food and Agriculture 2000; 80: 1033-1043.
29. Meydani M and Hasan ST: Dietary polyphenols and obesity. Nutrients 2010; 2: 737-751.
30. Vauzour D, Rodriguez-Mateos A, Corona G, Oruna-Concha MJ and Spencer JPE: Polyphenols and human health: prevention of disease and mechanisms of action. Nutrients 2010; 2: 1106-1131.
31. Shay J, Elbaz HA, Lee I, Zielske SP, Malek MH and Hüttemann M: Molecular mechanisms and therapeutic effects of (-)-Epicatechin and other polyphenols in cancer, inflammation, diabetes, and neurodegeneration. Oxidative Medicine and Cell Longevity 2015; ID 181260.
32. Hussain T, Tan B, Yin Y, Blachier F, Tossou MCB and Rahu. N: Oxidative stress and inflammation: what polyphenols can do for us? Oxidative Medicine and Cellular Longevity 2016; e7432797.

How to cite this article:

Attanayake AP: Dietary polyphenols: An overview. Int J Life Sci & Rev 2018; 4(11): 176-79. doi: 10.13040/IJPSR.0975-8232.IJLSR.4(11).176-79.

All © 2015 are reserved by International Journal of Life Sciences and Review. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)