



Received on 16 January, 2018; received in revised form, 04 February, 2018; accepted, 24 February, 2018; published 28 February, 2018

ANTIMICROBIAL ACTIVITY OF A SELECTED MEDICINAL PLANT EXTRACTS MIXTURE USED IN THE MANAGEMENT OF CHRONIC SKIN WOUNDS IN AYURVEDIC MEDICINE

W. G. G. Piumali¹, A. P. Attanayake^{*2} and H. Peiris¹

Department of Medical Laboratory Sciences¹, Faculty of Allied Health Sciences, University of Ruhuna, Karapitiya, Sri Lanka.

Department of Biochemistry², Faculty of Medicine, University of Ruhuna, Karapitiya, Sri Lanka.

ABSTRACT: The aim of the study was to determine the synergistic antimicrobial activity of a selected medicinal plant extract mixture composed of *Calotropis gigantean* (ElaWara, Family: Asclepiadaceae), *Datura metel* (Kalu Attana, Family: Solanaceae) and *Ricinus communis* (Erandu, Family: Euphorbiaceae) against selected Gram Positive and Gram Negative bacteria. In the study, antimicrobial activity was evaluated against two gram positive organisms; *Staphylococcus aureus* (ATCC 95923) and group A *Streptococcus spp.* (ATCC 12384) and two gram negative organisms; *Escherichia coli* (ATCC 95922) and *Pseudomonas aeruginosa* (ATCC 27853). Agar disc diffusion method was carried out to evaluate the antibacterial activity of the selected medicinal plant extract mixture and the inhibition zone diameters were measured. Further, minimum inhibitory concentration (MIC) was evaluated using broth micro-dilution method. The inhibition zones against *S. aureus* and group A *Streptococcus spp.* were observed. In contrast, inhibition zones were not demonstrated against selected gram negative organisms. The MIC evaluated for *S. aureus* was 16 mg/mL by broth dilution method. The results revealed that the selected medicinal plant extract mixture exerted synergistic antimicrobial activity against *S. aureus* and group A *Streptococcus spp.* However, antimicrobial activity against *E. coli* and *P. Aeruginosa* was not significant. Results were able to scrutinize the therapeutic benefits of the selected medicinal plant extracts mixture in the management of chronic skin wounds in Ayurvedic medicine. In addition, the plant extract mixture may offer valuable natural drug leads for the development of a commercially viable herbal antimicrobial formulation in future.

Keywords: Broth dilution method, Disc diffusion method, Medicinal plant extract mixture, Minimum inhibitory concentration, Sri Lankan Ayurveda medicine, Synergistic antimicrobial activity

Correspondence to Author:

Dr. A. P. Attanayake


Department of Biochemistry, Faculty of Medicine, University of Ruhuna, Karapitiya, Galle, Sri Lanka.

E-mail: anoja715@yahoo.com

INTRODUCTION: Throughout the history, man has had to contend with different kinds of skin wounds. Healing of chronic wounds, including pressure sores, leg ulcers, diabetic foot ulcers and other kinds of wounds, by secondary intention are common in community and clinical settings¹.

The management of chronic wounds represents a significant proportion of health-care resources and makes up a substantial amount of patient contact time in clinical setup.

The treatment of chronic wounds includes many strategies, including the use of various wound dressings, antimicrobial agents such as antibiotics, footwear, physical therapies and educational strategies. Use of antimicrobial agents for chronic wound care is promising than any other strategy^{2,3}. There is a wide range of antimicrobial agents with proven antimicrobial properties. In the past, acetic acid, chlorhexidine, honey, hydrogen peroxide,



sodium hypochlorite, potassium permanganate and proflavine have been used, however, the discovery and development of antibiotics during the 20th century has revolutionized the clinical therapy^{4, 5}. Antibiotics are one of the most important weapons in fighting bacterial infections. However, over the past few decades, these health benefits are under threat as many commonly used antibiotics have become less effective against certain conditions due to their toxic reactions and emergence of drug resistant bacteria⁶. The development of bacterial resistance is a huge problem since it is often not restricted to the specific antibiotic prescribed, but generally extends to other compounds of the same class⁷. Bacterial resistance and its rapid increase are the major concerns of global public health and are emerging as significant challenges to human health⁸.

However, the relentless emergence of antibiotic resistant strains of pathogens together with retarded discovery of novel antibiotics has led to discover antimicrobial agents specifically from herbal origin. In this sense, natural antimicrobials seem promising to many of the problems associated with the currently used antimicrobial agents in the management of wound infections. Unlike synthetic drugs, antimicrobials of plant origin are unlikely associated with side effects and exert a sufficient therapeutic potential to cure many infectious diseases^{9, 10}.

Finding healing powers in plants has a long history that began with folk medicine and through the years it has been incorporated into traditional and allopathic medicine. The mystique surrounding concepts in those traditional remedies has already led to an explosion of potential drug targets and numerous approaches have been developed to capture their intrinsic value. Among known plant species on the earth, only a small fraction has been investigated in the presence of antimicrobial compounds.

In fact, during the last few years, medicinal plants especially crude medicinal plant extract mixtures have attracted the attention of pharmaceutical and scientific communities, and evidence has demonstrated on promising potential of anti-microbial plant derived substances¹¹.

In Sri Lanka, Ayurvedic physicians use various plants in the management of wound healing as individual plants or as combinations of several plants. “Roganikash / Rohanekash” is a commonly used plant combination in the management of chronic skin wounds in Ayurvedic medicine. It consists of the leaves of *Calotropis gigantea* (ElaWara, Family: Asclepiadaceae), *Datura metel* (Kalu Attana, Family: Solanaceae) and *Ricinus communis* (Erandu, Family: Euphorbiaceae) (Fig. 1-3).



FIG. 1: CALOTROPIS GIGANTEA LINN.



FIG. 2: DATURA METEL LINN.



FIG. 3: RICINUS COMMUNIS LINN.

However, the therapeutic applications in the management of chronic skin wounds have not been scrutinized scientifically. The present study was to determine the synergistic antimicrobial activity of

MATERIALS AND METHODS:

Chemicals and Reagents: All chemicals and solvents were analytical grade and were used without purification.

Selection, Collection and Authentication of Plant Materials: The medicinal plants selected for the investigation are listed below. All fresh plant parts were collected from the Southern region of Sri Lanka. Botanical identity was determined by the descriptions given by Jayaweera (1982)¹². Voucher specimens were preserved at the Department of Medical Laboratory Science, Faculty of Allied Health Sciences, University of Ruhuna, Galle (FAHS/Piyumali/2017/1-3)

Extraction of Plants: The leaves of selected medicinal plants were cleaned and oven dried at 40°C until a constant weight was reached. Then, the dried plant materials were ground and the powder form of all plant material were mixed according to the formulae used in traditional medicine. Same amount of each plant powder was mixed together and dissolved in ethyl acetate. It was kept sealed in a conical flask and placed in a sonicator for 72 h. Then it was filtered and the filtrate was evaporated to dryness using a rotary evaporator. The percentage yield of the plant extract was 4.74 %.

Determination of Antimicrobial Activity:

Bacterial Strains, Culture Media and Growth Conditions: Antimicrobial effects were individually tested against four bacterial strains (two gram positive organisms; *Staphylococcus aureus* (*S. aureus*, ATCC 25923) and group A Streptococcus (ATCC 12384), two gram negative organisms; *Pseudomonas aeruginosa* (ATCC 27853) and *Escherichia coli* (ATCC 25922).

The standard cultures were used to evaluate antimicrobial activity of the plant extract mixture. The pure cultures were maintained on sheep blood agar medium for the entire study. Antimicrobial effect of the plant extract mixture was determined using agar disc diffusion method and broth dilution

the ethyl acetate extract of the above medicinal plant mixture composed of leaf extracts of *C. gigantean*, *D. metel* and *R. communis* against selected Gram Positive and Gram Negative bacteria.

method according to the Clinical and Laboratory Standards Institute (CLSI) protocols¹³. The suspensions of the tested micro-organisms were prepared with reference to the turbidity of the 0.5 McFarland standards.

Disc Diffusion Method: Filter paper discs were prepared from Whatman No.1 filter paper with a diameter of 6 mm. The discs were inserted into several small screw cap bottles and were autoclaved at 121 °C, 15 lb/in² for 15 minutes. A concentration of 1000 mg/mL of plant extract was prepared by dissolving the plant extract in DMSO (Dimethylsulfoxide) to obtain the concentration of 5000 µg/disc. Then, a graded dose of dilutions (400, 600 and 800 mg/mL) were prepared using the above stock solution to obtain the concentrations, 2000 µg/disc, 3000 µg/disc and 4000 µg/disc.

All solutions were well mixed using the vortex mixture. Filter paper discs were placed in sterile petri dishes with 5 cm distance from each disc. A volume of 5 µL of each dilution of plant extracts mixture and DMSO for negative controls were dispensed carefully on filter paper discs. Then petri dishes were kept closed for 30 minutes for drying of filter paper discs. Petri dishes containing Muller Hinton agar (MHA) and Muller Hinton Blood agar (MHA+5% human blood) for group A *Streptococcus spp.* were inoculated with relevant microbial suspensions. The filter paper discs impregnated with serial dilutions of plant extract solutions and with DMSO for negative controls were placed on the agar surface and were incubated over night at 37 °C. The diameter of any resulting zone of growth inhibition was measured¹³.

The clindamycin 2 µg and ciprofloxacin 5 µg were used as positive controls for gram positive and gram negative organisms respectively. Antimicrobial activity was determined by measuring the diameter of inhibition zone around the disc.

Broth Micro-dilution Method: The minimum inhibitory concentration (MIC) was determined for the organisms (except group A Streptococcus) which were sensitive to the plant extract mixture

previously evaluated by the broth dilution method according to the guidelines of CLSI¹³. Serially diluted plant extract was added to sterile 96-well microtiter plates which contain bacterial suspension and Muller Hinton broth growth medium according to the standard CLSI protocol¹³. Muller Hinton broth was prepared and aliquoted into khan tubes. The broth volumes were prepared (1490 μ L, 450 μ L and 2.5 mL) and autoclaved.

A graded concentration of plant extract solutions, 480 mg/mL, 440 mg/mL, 400 mg/mL, 360 mg/mL, 320 mg/mL and 280 mg/mL were prepared. A volume of 50 μ L of plant extract solution is mixed with 450 μ L of Muller Hinton broth which was previously prepared in khan tubes. It was well mixed and 100 μ L of the resultant was transferred to the first well of the microtiter plate. A volume of 50 μ L of Muller Hinton Broth was added to the rest of the wells. Then 50 μ L of first well was transferred to the second and well mixed again.

A volume of 50 μ L was transferred from the second to the third. The same procedure was repeated except in the last well which was considered as the negative control. Then a volume of 10 μ L of organism suspension, which was adjusted to the turbidity of 0.5 McFarland, was added to 1490 μ L of Muller Hinton broth to obtain the final inoculum size 10⁶ CFU/mL. It was well mixed and 50 μ L was added to each well except the well before last well.

The last well was remained un-inoculated, to identify the wells with complete growth inhibitions of the organism by comparing with it. The plate was mildly shaken carefully and was incubated at 37°C for 24 h. The minimum inhibitory concentration (MIC) is defined as the lowest concentration of antimicrobial agent that completely inhibits the growth of organisms in the micro-dilution wells as detected by the unaided eye.

The presence and inhibition of micro-organisms were identified by the turbidity in the broth. All dilutions in the wells were inoculated on a growth media to confirm the inhibition of the organism by the plant extract. Several concentrations of the plant extract mixture were evaluated for the MIC until success¹⁴.

Statistical Analysis: Each test was performed in triplicates. Data will be expressed as mean \pm standard deviation (SD).

RESULTS AND DISCUSSION: In the present study, the antimicrobial activity of a selected medicinal plant extract mixture which has been used as an antimicrobial medicine by Ayurvedic physicians was evaluated. The activity was tested against, wound associated pathogens; *S. aureus*, group A *Streptococcus spp.*, *E. coli* and *P. aeruginosa*. The plant combination was consists of three medicinal plants namely *C. gigantea*, *D. metel* and *R. communis*. The selected medicinal plants parts were collected during July to August 2017, from their natural habitat.

Antimicrobial properties of medicinal plants are being increasingly reported from different parts of the world. Accordingly a number of single medicinal plants grown in Sri Lanka were subjected to determine the antimicrobial activity, however very few medicinal plant extract mixtures were used to determine the antimicrobial activity. However, the antimicrobial activity of the selected medicinal plant extract mixture was not investigated previously. This is the first report on scrutinization of the selected herbal mixture towards it's antimicrobial activity.

The single medicinal plant extract of *C. gigantea* showed antibacterial activity against number of organisms. The antibacterial activity of the methanol, petroleum ether, chloroform and ethyl acetate extract from the root bark of *C. gigantean* and its sub fractions were investigated using agar disc diffusion method. Its root bark extracts was active against *P. aeruginosa* and *E. coli* at 20, 30 and 40 μ g/disc doses. The extracts showed inhibition zones more than 15 mm of diameter for both *P. aeruginosa* and *E. coli*¹⁵. In the present study, as shown in **Fig. 4, 5, 6, 7, 8**, there were no inhibitory zones for *S. aureus* *Pseudomonas aeruginosa* and *E. coli*. It implied that the individual plant extract of *C. gigantea* exerted a significant antimicrobial potential than that of in its combination. Another study showed that the crude extract of leaves of the *C. gigantea* exerted antimicrobial activity against *S. aureus*, *P. aeruginosa* and *E. coli*.

According to the previous reports, the MIC was evaluated by modified agar well diffusion method and the MIC values for *S. aureus*, *E. coli* and *P. aeruginosa* were 50 mg/mL, 12.5 mg/mL, 3.1 mg/mL respectively. In the present study there was no activity against *E. coli* and *P. aeruginosa* and 16 mg/mL of MIC was determined for *S. aureus* (Table 1, 2, 3 and Fig. 5). However, the results showed that the plant combination including the *C. gigantea* is more effective against *S. aureus* than in the single extract of *C. gigantea*. The antimicrobial activity of leaf, stem bark and root extracts of *D. metel* was evaluated by agar well diffusion method,

against β hemolytic Streptococcus, *E. coli*, *P. aeruginosa* and *S. aureus*. They were sensitive to the ethanol and aqueous leaf and stem bark extracts of *D. metel*. The root extracts of the plants had no antibacterial activity. The leaf extracts exerted potential effects on above isolates. The crude ethanol extract exhibited an inhibitory zone of more than 30 mm against *P. aeruginosa*. It was found that the inhibitory zones were more than 20 mm for *E. coli*, *S. aureus*, and β haemolytic Streptococcus according to previous reports. The MIC evaluated using broth macro dilution method for these organisms were 20 mg/mL¹⁶.



FIG. 4: ZONES OF INHIBITION AT VARIOUS CONCENTRATIONS OF PLANT EXTRACT MIXTURE AGAINST *S. AUREUS* IN DISC DIFFUSION METHOD




FIG. 5: ZONES OF INHIBITION AT VARIOUS CONCENTRATIONS OF PLANT EXTRACT MIXTURE AGAINST GROUP A STREPTOCOCCUS IN DISC DIFFUSION METHOD




FIG. 6: ZONES OF INHIBITION AT VARIOUS CONCENTRATIONS OF PLANT EXTRACT AGAINST *E. COLI* IN DISC DIFFUSION METHOD




FIG. 7: ZONES OF INHIBITION AT VARIOUS CONCENTRATIONS OF PLANT EXTRACT AGAINST *P. AERUGINOSA* IN DISC DIFFUSION METHOD

TABLE 1: INHIBITORY ZONE DIAMETERS OBTAINED IN DISC DIFFUSION METHOD

Plant concentration ($\mu\text{g}/\text{disc}$)	Diameter of the inhibitory zone, M \pm SD (mm)				
	2000	3000	4000	5000	Pos. control
<i>Staphylococcus aureus</i>	7.7 \pm 0.6	7.7 \pm 0.6	8.7 \pm 0.6	8.7 \pm 0.6	27.7 \pm 1.5
Group A Streptococcus	8.3 \pm 0.6	8.7 \pm 0.6	9.7 \pm 0.6	10.0 \pm 1.0	32.7 \pm 1.2
<i>Escherichia coli</i>	-	-	-	-	32.7 \pm 0.6
<i>Pseudomonas aeruginosa</i>	-	-	-	-	32.0 \pm 0.0

M \pm SD: mean of the zone diameter \pm standard deviation

In the present study, inhibitory zones were not observed for *E. coli* and *P. aeruginosa*. As shown in **Table 2**, the plant extract showed an inhibitory zone at a diameter of (7.6 ± 0.6 , 7.8 ± 0.6 , 8.7 ± 0.6 , 8.7 ± 0.6) mm for the concentrations, 400, 600, 800, 1000 mg/mL against *S. aureus* respectively. The deviations in results of the present investigation may be due to the difference of method used to evaluate antimicrobial activity, method of extraction, extraction solvent and the difference in phytochemicals present in the plant extracts.

In disc diffusion method, small quantity of plant extract is saturated in a disc and inhibitory zone surrounding the disc was observed. The MIC evaluated for *S. aureus* in the present study was 16 mg/mL. The value was quite smaller than the reported values in previous studies.

It showed that the leaf extract of *D. metel* in combination is more effective than the individual leaf extract of it, against *S. aureus*. The antimicrobial activity was evaluated by agar well diffusion method in ethanol and ethyl acetate extracts of mature leaves of *D. metel* against *S. aureus*, *E. coli* and *P. aeruginosa*.

The ethyl acetate extract of *D. metel* (100 mg/mL) showed the maximum zone of inhibition of 19 mm against *E. coli*. There was no zone of inhibition demonstrated against *P. aeruginosa*. Inhibitory zone (13 mm diameter) was observed for *S. aureus*. The ethanol extract of the same plant, *D. metel* (100 mg/mL) showed the maximum zone of inhibition (26 mm) against *P. aeruginosa* and *E. coli*. *S. aureus* showed a relatively minor zone of inhibition (8 mm). This implied that the antimicrobial activity of the plant extracts depend on the solvent used for the extraction process¹⁷.

TABLE 2: MIC DETERMINATION USING THE PLANT EXTRACT AT THE CONCENTRATION OF 480 mg/mL

Concentration of plant extract in contact with bacteria in microtiter wells (mg/mL)	Clear or turbid (+/-)
48	+
24	+
12	-
6	-
3	-
0	-

(-): Clear, (+): Turbid

TABLE 3: DETERMINATION OF MINIMUM INHIBITORY CONCENTRATION WITH TWO FOLD DILUTIONS

Original concentration of the plant extract solution (mg/mL)	Concentration of plant extract in contact with bacteria in microtiter wells (mg/mL)			
440	44	22	11	5.5
400	40	20	10	5.0
360	36	18	9	4.5
320	32	16	8	4.5
280	28	14	7	3.5

In the present investigation, at the concentrations of 400, 600, 800, 1000 mg/mL, *S. aureus* exhibited inhibitory zones with (7.6 ± 0.6 , 7.8 ± 0.6 , 8.7 ± 0.6 , 8.7 ± 0.6) mm of diameters respectively (**Table 2**). The inhibitory zones were not obtained for *E. coli* and *P. aeruginosa*. However, the plant extract mixture was more powerful against *S. aureus*. The individual plant, *D. metel* was more active against *E. coli* than its' in the combination.




FIG. 8: MIC DETERMINATION IN A MICROTITER PLATE USING BROTH MICRO-DILUTION METHOD MIC OF *S. AUREUS* WAS 16 mg/mL

Methanol, ethanol and aqueous leaf extracts of *R. communis* had antimicrobial potential against *S. aureus* and *P. aeruginosa*. The inhibitory zones with diameters between 15 to 20 mm were obtained against *S. aureus* and *P. Aeruginosa* by agar well diffusion method¹⁸.

In the present study, no inhibitory zones were obtained against *P. aeruginosa* and *E. coli*. The results could not be compared straightway since the antimicrobial activity was evaluated using two different methods as agar well diffusion and agar disc diffusion methods. However, the individual plant, *R. communis* as well as in combination is active against *S. aureus*. The individual plant is more effective against gram negative bacterial organisms than in its' combination.

While many studies have focused on the antimicrobial activity of individual plants, only few have reported on plant - plant interactions when used in combination. Synergistic interactions noted for *Pentanisia prunelloides* combined with *Elephantorrhiza elephantina* validate an enhanced antimicrobial effect when used in combination¹⁹. Thripala which is commonly used Sri Lankan formulation of three medicinal plants, namely, *Terminalia chebula*, *Terminalia bellerica* and *Embllica officinalis* were subjected for the synergistic antimicrobial activity against common pathogens^{20, 21}.

The antimicrobial activity of essential oil of *Laurus nobilis* L. and *Myrtus communis* L. in combination have shown a great potential with mild heat and high hydrostatic pressure to obtain a higher inactivation of food borne pathogens²². According to the results of previous studies, individual medicinal plants among the plants selected in the present study, are potentially active against microbes than they are in combination. The leaf extract of *D. metel* is less susceptible to gram negatives especially *P. aeruginosa*.

However, *C. gigantea* and *R. communis* are susceptible to gram positive pathogens; *S. aureus* and group A *Streptococcus*, as well as gram negative pathogens; *E. coli* and *P. aeruginosa*. So the activity of individual plants could be more susceptible than they were in combination. It is difficult to compare the results because most of the studies have done using agar well diffusion method in which a well contains large quantity of the plant extract than in a disc diffusion method.

However, the evaluation of antimicrobial activity of individual plants as well as their combination, using the same methodology is important in interpretation and comparison of antimicrobial activity more precisely. Ethyl acetate extract was used in the present study due to its high activity against microorganisms²². The medicinal properties of plant extracts normally depend upon the presence of active compounds possessing specific functional groups that are soluble only in solvents of particular polarity. Studies have shown that ethyl acetate extracts contain potent active compounds against microbes²².

There was a significant difference of inhibitory zone diameters with the reference antibiotic. In the present study, clindamycin 2 µg and ciprofloxacin 30 µg were selected as the positive controls against gram positive pathogens and gram negative pathogens respectively. Clindamycin is a lincosamide antibiotic that is active against many aerobic Gram-Positive cocci and a range of anaerobic Gram-Positive and Gram-Negative bacteria. Ciprofloxacin is a fluoroquinolone antibiotic, with a broad spectrum of activity targeting aerobic Gram-Negative pathogens²³. The diameters of inhibitory zones obtained positive controls were obviously higher than the values obtained for antimicrobial discs prepared from the plant extract. It could be due to the fact that the active compound was only a small percentage of the extract since no purification was done at this stage.

CONCLUSION: The ethyl acetate extract of the selected medicinal plant extract mixture composed of the leaves of *Calotropis gigantea*, *Datura metel* and *Ricinus communis*, showed a significant antimicrobial activity against gram positive pathogens, *Staphylococcus aureus* and group A streptococcus spp. in agar disc diffusion method. The minimum inhibitory concentration for *S. aureus* was 16 mg/mL in broth micro-dilution method. However, the plant extract mixture had no activity against the gram negative pathogens, *Escherichia coli* and *Pseudomonas aeruginosa*. Results were able to scrutinize the therapeutic benefits of the selected medicinal plant extract mixture in the management of chronic skin wounds in Ayurvedic medicine. In addition, the plant extract mixture may offer valuable natural drug leads for the development of a commercially viable herbal antimicrobial formulation.

ACKNOWLEDGEMENT: Nil

CONFLICTS OF INTEREST: Nil

REFERENCES:

1. Sun X, Jiang K, Chen J, Wu L, Lu H, Wang A and Wang J: A systematic review of maggot debridement therapy for chronically infected wounds and ulcers. International Journal of Infectious Diseases 2014; 25: 32-37.
2. Cooper R: A review of the evidence for the use of topical antimicrobial agents in wound care. World Wide Wounds 2004; <http://www.worldwidewounds.com/2004/february/Cooper/Topical-Antimicrobial-Agents.html>

3. O'Meara S, Cullum N, Majid M and Sheldon T: Systematic reviews of wound care management: (3) antimicrobial agents for chronic wounds; (4) diabetic foot ulceration. *Health Technology Assessment* 2004; 4(21): 1-237.
4. Boucher HW, Talbot GH, Bradley JS, Edwards JE and Gilbert D: Bad bugs, no drugs: No escape! An update from the infectious diseases society of America. *Clinical Infectious Diseases* 2009; 48: 1-12.
5. Pawar PL and Nabar BM: Effect of plant extracts formulated in different ointment bases on MDR Strains. *Indian Journal of Pharmaceutical Sciences* 2010; 72(3): 397-401.
6. Aiyegoro OA, Afolayan AJ and Okoh AI: Synergistic interaction of *Helichrysum pedunculatum* leaf extracts with antibiotics against wound infection associated bacteria. *Biological Research* 2009; 42(3): 327-338.
7. Levy SB and Marshall B: Antibacterial resistance worldwide: causes, challenges and responses. *Nature Medicine* 2004; 10: 122-129.
8. Aiyegoro OA, Afolayan AJ and Okoh AI: *In-vitro* antibacterial activities of crude extracts of the leaves of *Helichrysum longifolium* in combination with selected antibiotics. *African Journal of Pharmacy and Pharmacology* 2009; 3(6): 293-300.
9. Chanda S, Dudhatra S and Kaneria M: Antioxidative and antibacterial effects of seeds and fruit rind of nutraceutical plants belonging to the family Fabaceae family. *Food and Function* 2010; 1: 308-315.
10. Habbal O, Hasson SS, El-Hag AH, Al-Mahrooqi Z, Al-Hashmi N, Al-Bimani Z, Al-Balushi, MS and Al-Jabri AA: Antibacterial activity of *Lawsonia inermis* Linn (Henna) against *Pseudomonas aeruginosa*. *Asian Pacific Journal of Tropical Biomedicine* 2011; 1: 173-176.
11. Misra LN, Wouatsa VNA, Kumar S, Kumar VR and Tchoumboungang F: Antibacterial, cytotoxic activities and chemical composition of fruits of two *Cameroonian zanthoxylum* species. *Journal of Ethno-pharmacology* 2013; 148(1): 74-80.
12. Jayaweera DMA: Medicinal plants (indigenous and exotic) used in Ceylon. Sri Lanka: National Science Foundation in Sri Lanka, 1982.
13. CLSI: Clinical and laboratory standards institute catalog 2014.
14. Balouiri M, Sadiki M and Ibsouda SK: Methods for *in vitro* evaluating antimicrobial activity: A review. *Journal of Pharmaceutical Analysis* 2016; 6(2): 71-79.
15. Alam MA, Habib MR, Nikkon R, Rahman M and Karim MR: Antimicrobial activity of Akanda (*Calotropis gigantea* L.) on some pathogenic bacteria. *Bangladesh Journal of Scientific and Industrial Research* 2008; 43(3): 397-404.
16. Akharaiyi FC: Antibacterial, phytochemical and antioxidant activities of *Datura metel*. *International Journal of Pharm Tech Research*. 2011; 3(1): 478-483.
17. Saranraj P: Antibacterial evaluation and phytochemical screening of *Datura metel* leaf extracts against bacterial pathogens. *International Journal of Pharmaceutical and Biological Archive* 2011; 2(4): 1130-1136.
18. Naz R and Bano A: Antimicrobial potential of *Ricinus communis* leaf extracts indifferent solvents against pathogenic bacterial and fungal strains. *Asian Pacific Journal of Tropical Biomedicine*. 2012; 2(12): 944-949.
19. Safiullah A, Harish CC, Vijay AK and Saira K: Antimicrobial activity of Triphala against bacterial isolates from HIV infected patients. Jundishapur. *Journal Microbiology* 2011; 4(1): 9-17.
20. Sheta MM, Kikani KM, Kavathia P, Thakkar J and Rangnani T: Study of antimicrobial activity of Triphala and its individual components *International Journal of Herbal Medicine*. 2016; 4(5): 41-43.
21. Cherrat L, Espina L, Bakkali M, García-Gonzalo D, Pagán R and Laglaoui A: Chemical composition and antioxidant properties of *Laurus nobilis* L. and *Myrtus communis* L. essential oils from Morocco and evaluation of their antimicrobial activity acting alone or in combined processes for food preservation. *Journal of the Science of Food and Agriculture* 2014; 94(6): 1197-1204.
22. Martin WW, Matasyoh JC and Kinyanjui T: Antimicrobial activity of solvent extracts from the leaves of *Tarchonanthus camphoratus* (Asteraceae). *Journal of Pharmacognosy and Phytochemistry* 2014; 3(1): 123-127.
23. Card RM, Mafura M, Hunt T, Kirchner M, Weile J, Rashid MU, Weintraub A, Nord CE and Anjum MF: Impact of ciprofloxacin and clindamycin administration on Gram-Negative bacteria isolated from healthy volunteers and characterization of the resistance genes they harbor. *Antimicrobial Agents and Chemotherapy* 2015; 59(8): 4410-4416.

How to cite this article:

Piumali WGG, Attanayake AP and Peiris H: Antimicrobial activity of a selected medicinal plant extracts mixture used in the management of chronic skin wounds in Ayurvedic medicine. *Int J Life Sci Rev*. 2018; 4(2): 27-34. doi: 10.13040/IJPSR.0975-8232.IJLSR.4(2).27-34.

All © 2015 are reserved by International Journal of Life Sciences and Review. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to ANDROID OS based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)