

Received on 15 April 2016; received in revised form, 16 May 2016; accepted, 19 May 2016; published 31 May 2016

IN-VITRO ANTI-HAEMOLYTIC, ANTI-BACTERIAL AND PHYTO-CONSTITUENTS INVESTIGATION OF *ANDROGRAPHIS PANICULATA* METHANOLIC LEAF EXTRACT

Venkata Narasimha Kadali ^{*1}, Tadi Ramesh ², Sudhakara Rao Pola ¹ and B. V. Sandeep ¹

Department of Biotechnology ¹, Andhra University, Visakhapatnam - 530003, Andhra Pradesh, India.

Department of Biotechnology ², S V K P & Dr. K.S. Raju Arts and Science College, Penugonda - 534320, Andhra Pradesh, India.

ABSTRACT: The bitter plant *Andrographis paniculata* has been selected to evaluate the anti haemolytic activity, antibacterial activity, and various phytochemicals present in the methanolic leaf extract. The leaf extract of *Andrographis paniculata* shown to have significant anti-haemolytic activity. The extract showed anti-haemolytic activity in the range from 42% to 80%. At varying concentration of plant extracts 20, 40, 60, 80 and 100 mg/ml the percent of inhibition of haemolysis recorded were 42%, 56%, 62%, 70%, 80% respectively. The methanolic leaf extract of *Andrographis paniculata* showed significant inhibitory effect on all of the five bacterial species such as *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Escherichia coli*, *Bacillus subtilis*, and *Streptococcus mutans*. The inhibition zones for *S. aureus* were 12 mm, 14 mm and 19 mm (including well 4 mm) at concentrations of 15 µl, 20 µl, and 25 µl respectively. The inhibition zones *P. aeruginosa* were 11 mm, 13 mm and 20 mm, 8 mm, 10 mm and 15 mm for *Streptococcus mutans*. The zones for *E. coli* were 15 mm, 16 mm and 18 mm and 12 mm, 13 mm and 19 mm for *Bacillus subtilis*. The phytochemical investigation of methanolic extract of *Andrographis paniculata* leaves showed the presence of carbohydrates, flavonoids, alkaloids, tannins, phenols, and saponins.

Keywords: *Andrographis paniculata*, Phytochemicals, Anti-haemolytic, Anti-bacterial

Correspondence to Author:

Venkata Narasimha Kadali

Department of Biotechnology, Andhra University, Visakhapatnam - 530003, Andhra Pradesh, India.

E-mail: vnsimhakadalibiotech@gmail.com

INTRODUCTION: Despite the remarkable progress in the preparation of synthetic drugs, over 25% of prescribed medicines in industrialized countries are derived directly from plants ¹. The World Health Organization (WHO) also considered phytotherapy in its health programs and suggested basic procedures for validation of drugs from plant origin in developing countries ².

Indigenous plants are reservoirs of various metabolites and provide an unlimited source of important chemicals that have diverse biological properties ³. Medicinal plants are gaining a lot of importance nowadays because of efficacy they have been showing in the traditional healing ⁴. Herbs are the source of magnificent inhibitors that could act on a wide variety of diseases. One of the great aspects of herbs is they show 100% results when it comes to the healing. Herbs have all sorts of answers against various diseases ⁵.

The best source of drugs without lethal effects to human systems could be the plant source, and this has been proved by the Traditional healing system and the recent studies conducted on the

<p>QUICK RESPONSE CODE</p> 	<p>DOI: 10.13040/IJPSR.0975-8232.IJLSR.2(5).102-08</p> <p>The article can be accessed online on www.ijlsr.com</p>
<p>DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.IJLSR.2(5).102-08</p>	

experimental animals⁶. Even expansion of modern medicines throughout the India people in rural areas still uses this wonder herbal medications for various sorts of diseases⁷. The use of medicinal plants in developing countries is increasing, which offer a new source of antibacterial, antifungal, and antioxidants agents⁸. The development of drug resistance, as well as the appearance of undesirable side effects of certain drugs, has led to the search of new antimicrobial agents in particular from medicinal plants⁸. This medicinal herb *A. paniculata* otherwise called as king of bitters because it has extremely bitter where it is used to treat various infections and diseases⁹.

Occurs quite commonly in the entire India subcontinent, which accounts for its widespread use since ancient times against a variety of disorders, in both Ayurvedic and Chinese Medicine¹⁰. It has blood purifying property, so it is recommended for use in leprosy, gonorrhoea, scabies, boils, skin eruptions, and chronic and seasonal fevers⁹. The leaf extract of *Andrographis paniculata* was used to evaluate the enzyme inhibiting the activity of protease, phosphomonoesterase, phosphodiesterase, acetylcholinesterase, phospholipase A2, hyaluronidase and L-amino acid oxidase toxic enzymes present in snake venom¹¹.

The purified active compounds through TLC and crude plant extract were subjected to analyze the anti-inflammatory activity and shows potent inhibitory activity against the adverse effects of the inflammation such as tissue damage and protein denaturation¹⁰. *A. paniculata* shows excellent anticancer activities against different cancer cell lines; it is alternatives medicines for cancer would replace side effect causing chemotherapeutic agent¹².

Andrographolide active component of *A. paniculata* shown analgesic, antipyretic, and anti-ulcerogenic effect¹³. Anti-hyperglycemic effect in streptozotocin-induced diabetic rats¹⁴, it has anti-tumor activity via activation of cytotoxic T lymphocyte and attenuation of tumor growth *in-vivo*¹⁵. *A. paniculata* is known to possess anti-HIV¹⁶, cardioprotective¹⁷ and hepatoprotective¹⁸, properties. In this present study, the bitter plant *Andrographis paniculata* has been selected to evaluate the anti haemolytic activity, antibacterial

activity, and various phytochemicals present in the methanolic leaf extract.

MATERIALS AND METHODS:

Materials: Leaves of *Andrographis paniculata* were collected from the campus S V K P & Dr. K S Raju Arts and Science College, Penugonda. The plant was authenticated by Dr. Suryanayana Raju, Department of Botany, S V K P & Dr. K S Raju Arts and Science College, Penugonda. Leaves were dried in sunlight for a week and then powdered using a blender to get a coarse powder. Chicken blood was collected from the local market in EDTA bottle.

Test Microorganisms: The methanolic leaf extract of *Andrographis paniculata* was tested against five pathogenic bacteria. The test organisms include *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Escherichia coli*, *Bacillus subtilis*, and *Streptococcus mutans*.

Extraction Process: Preparation of extract was done according to the procedure done by Sharmila et al., (2011)¹⁹. 25g of leaf powder was packed in a Soxhlet extraction unit and exhaustively extracted using 100 ml of methanol at 60 °C for 12 h. The extract was completely dried in a water bath at 40 °C and subsequently stored at 4 °C.

Phyto-chemical Analysis: A preliminary phyto-chemical investigation was conducted for the detection of steroids, terpenoids, flavonoids, saponins, tannins, carbohydrates, and phenols by using standard procedures^{20,21}.

Determination of Antibacterial Activity: Antibacterial activity was measured by well diffusion method²². Nutrient agar (Hi-media) was prepared and poured into the Petri plates. After solidification of media, overnight bacterial cultures were inoculated on the surface of media. By using a sterile gel puncher, 4 mm of wells were made in each Petri plates.

Then 15 µl, 20 µl, 25 µl of methanolic leaf extract *Andrographis paniculata* were added into the three wells respectively. The plates were incubated in the incubator at 37°C for optimum bacterial growth. In the next day, the diameter of the zone of inhibition was measured.

Assessment of Anti-haemolytic Activity: Anti-haemolytic activity against H₂O₂ induced haemolysis in chicken RBC determined by *in-vitro* method described by Tavazzi et al., 2001, Thagriki Dluya et al., 2015^{23, 24}. The chicken erythrocytes were separated by centrifugation at 200 rpm and washed with saline or isotonic sodium phosphate buffer (pH 7.4) until the supernatant is colorless. The erythrocytes were then diluted with saline or phosphate buffer to give a 4% suspension. Varying amounts of the plant extracts (20, 40, 60, 80 and 100 mg /ml) with saline or buffer was added to 2 ml of the suspension of erythrocytes, and the volume was made up to 3.5 ml with saline or buffer. This mixture was pre-incubated for 120 min and then 0.5 ml H₂O₂ solutions of appropriate concentration in saline or buffer was added. The concentration of H₂O₂ in the reaction mixture was adjusted to bring 90% haemolysis of blood cells after 120 min incubation. After 120 min of

incubation, tubes were centrifuged, and the amount of haemolysis was determined by measurement of the absorbance at 540 nm corresponding to haemoglobin liberation. The anti-haemolytic activity was expressed as the inhibition percentage and was calculated using the following formula:

$$\text{Anti-haemolytic activity (\%)} = \frac{\text{Control 540 nm} - \text{Sample 540 nm}}{\text{Control 540 nm}} \times 100$$

Where, Sample 540 nm was the absorbance of the sample and Control 540 nm was the absorbance of the control.

RESULTS:

Phytochemical Analysis: The phytochemical investigation of methanolic extract of *Andrographis paniculata* leaves showed the presence of carbohydrates, flavonoids, alkaloids, tannins, phenols, and saponins **Table 1**.

TABLE 1: PRESENCE OF PHYTOCHEMICALS IN METHANOLIC EXTRACT OF ANDROGRAPHIS PANICULATA LEAVES

S. no.	Phytoconstituent	Presence of phytochemical in methanolic extract
1	Carbohydrates	+
2	Flavonoids	+
3	Alkaloids	+
4	Tannins	+
5	Phenols	+
6	Proteins	-
7	Saponins	+
8	Sterols	-

+ indicates presence; - indicates absence.

Antibacterial Activity: The methanolic leaf extract of *Andrographis paniculata* showed significant inhibitory effect on all of the five bacterial species such as *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Escherichia coli*, *Bacillus subtilis*, and *Streptococcus mutans*. The inhibition zones for *S. aureus* were 12 mm, 14 mm and 19 mm (including well 4 mm) at concentrations of 15

µl, 20 µl, and 25 µl respectively. The inhibition zones *P. aeruginosa* were 11 mm, 13 mm and 20 mm, 8 mm, 10 mm and 15 mm for *Streptococcus mutans*. The zones for *E. coli* were 15 mm, 16 mm and 18 mm and 12 mm, 13 mm and 19 mm for *Bacillus subtilis*. The antibacterial results of methanolic extract were given below **Table 2** and **Fig. 1-6**.

TABLE 2: INHIBITION ZONES OF METHANOLIC EXTRACT OF ANDROGRAPHIS PANICULATA AGAINST TEST PATHOGENIC BACTERIA

Test bacterial species	Inhibition zones in mm (including well 4 mm)		
	15 µl (methanolic extract)	20 µl (methanolic extract)	25 µl (methanolic extract)
<i>Staphylococcus aureus</i>	12 mm	14 mm	19 mm
<i>Pseudomonas aeruginosa</i>	11 mm	13 mm	20 mm
<i>Bacillus subtilis</i>	12 mm	13 mm	19mm
<i>Escherichia coli</i>	15 mm	16 mm	18 mm
<i>Streptococcus mutans</i>	8 mm	10 mm	15 mm



FIG. 1: STAPHYLOCOCCUS AUREUS



FIG. 2: STREPTOCOCCUS MUTANS



FIG. 3: ESCHERICHIA COLI



FIG. 4: BACILLUS SUBTILIS



FIG. 5: PSEUDOMONAS AERUGINOSA




FIG. 6: ZONES OF INHIBITION OF METHANOLIC EXTRACT ON DIFFERENT BACTERIAL SPECIES

Anti-haemolytic Activity: The methanolic leaf extract of *Andrographis paniculata* shown to have significant anti-haemolytic activity. The methanolic extract showed anti-haemolytic activity in the range

from 42% to 80%. At varying concentration of plant extracts 20, 40, 60, 80 and 100 mg/ml the percent of inhibition of haemolysis recorded were 42%, 56%, 62%, 70%, 80% respectively **Fig. 7**.




FIG. 7: ANTI-HAEMOLYTIC ACTIVITY OF METHANOLIC LEAF EXTRACT OF ANDROGRAPHIS PANICULATA

DISCUSSION: The preliminary phytochemical investigation of methanolic extract of *Andrographis paniculata* leaves showed the presence of carbohydrates, flavonoids, alkaloids, Tannins, phenols, and saponins. The methanolic leaf extract of *Andrographis paniculata* shown to have significant anti-haemolytic activity. In this study, the inhibition of haemolysis found to be increased with increase in the concentration of extract. When red blood cells were treated with H_2O_2 (toxicant), % haemolysis was found to be increased. This may be because of the oxidizing nature of H_2O_2 concerning cell membrane degradation and release of haemoglobin from the cell²⁵. H_2O_2 also cause mobilization of Fe^{2+} by Ca^{2+} via Fenton reduction stimulating the production of OH- radicals²⁶.

All these factors combinedly cause destabilization of the cell membrane, which is probably the key event of the lysis of the cell²⁵. Polyphenols possess many biological effects, mainly attributed to their antioxidant activities in scavenging free radicals, inhibition of peroxidation and chelation of transition metals^{27, 28}. Chidambaram et al., (2011)²⁹ proved that *Andrographis paniculata* has significant ability to stabilize the Red blood cell membrane. These findings are by them. The methanolic leaf extract of *Andrographis paniculata*

showed inhibitory effect on all of the five bacterial species such as *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Escherichia coli*, *Bacillus subtilis*, and *Streptococcus mutans*. It is shown that methanol extract had a broad spectrum of activity. The extract has shown inhibitory zones ranging from 8 mm to 20 mm.

In all the bacteria used in this study, *Pseudomonas aeruginosa* proved to be more susceptible and exhibited maximum inhibitory zone 20 mm. The minimum inhibitory zone was observed for *Streptococcus mutans* 15 mm. As modern antibiotics have various anarchic toxic effects, these plant extracts could serve as an alternative antibacterial agent. In single plant many active secondary metabolites are present and medicinal effect can be attributed to either to a single compound or synergistic effect of many compounds and present antibacterial activity can either be due to the presence of some specific bioactive molecule or due to the synergistic effect of different phytoconstituents³⁰.

CONCLUSION: Through this present study, it can be concluded that the methanolic extract of *A. paniculata* leaves showed significant antibacterial activity and membrane stabilizing activity. These effects are due to the phyto inhibitors present in the

leaf extract. The feasibility of making efficient drugs from plant sources is very much near. Further research should be done to isolate inhibitors and necessarily formulated.

ACKNOWLEDGEMENT: The authors wish to thank the secretary and correspondent of S.V.K.P & Dr. K. S. Raju Arts & Science College, Penugonda for providing the research laboratory and also chemicals which enabled us to complete this work. The authors are grateful to the expertise available in the campus that helped a lot in species identification.

CONFLICT OF INTEREST: Nil

REFERENCES:

- Newman DJ, Cragg GM and Snader KM: The influence of natural products upon drug discovery. *Natural Product Reports* 2000; 17(3): 215-234.
- Anushia C, Sampathkumar P and Ramkumar L: Antibacterial and antioxidant activities in *Cassia auriculata*. *Global Journal of Pharmacology* 2009; 3(3): 127-130.
- Tomoko N, Takashi A, Hiromi T, Yuka I, Hiroko M, Munekazu I, Totshiyuki T, Tetsuro I, Fujio A, Iriya I, Tsotomu N and Kazuhito W: Antibacterial activity of extracts prepared from tropical and subtropical plants on methicillin-resistant *Staphylococcus aureus*. *Journal of Health Sciences* 2002; 48: 273-276.
- Kadali VN and Sandeep BV: Anti-hyperglycemic plants used by the traditional healer of west Godavari District, Andhra Pradesh, India. *Int J Pharmacognosy* 2015; 2(9): 473-77. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.2\(9\).473-77](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.2(9).473-77).
- Kadali VN, Kindangi KR, Rao PS and Sandeep BV: Wonder herbs having anti asthmatic activity present in West Godavari District, Andhra Pradesh, India- A Mini Review. *Advances in Biology, Biotechnology and Genetics* 2016; 3(1): 01-06.
- Kadali VN, Kindangi KR, Peter AE, Sudhakara Rao P, Bindiya P and Sandeep BV: Hepato-Protective Herbs-Present in West Godavari District of Andhra Pradesh, India- A mini review. *International Journal of Medical and Health Research* 2015; 1(1): 15-18.
- Kadali VN, Kindangi KR, Pola SR, Ramesh T and Sandeep BV: Cardio-protective plants present in West Godavari District of Andhra Pradesh, India: A Short Review. *Int J Pharmacognosy* 2016; 3(1): 100-07. doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.3\(1\).100-07](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.3(1).100-07).
- Premanath R and Devi NL: Antibacterial, antifungal and antioxidant activities of *Andrographis paniculata* Nees. *Leaves. IJPSR* 2011; 2(8): 2091-2099.
- Rajeshkumar S: Antimicrobial effect of King of bitter *Andrographis paniculata* and traditional herb *Aegle marmelos* against clinical pathogens. *Int J Pharm Tech Res* 2014-2015; 7(2): 325-329.
- Sundar S, Sudevan S, Paramasivam R and Vijayaraghavan R: Evaluation of *in-vitro* anti-inflammatory activity of aqueous extract of *Andrographis paniculata*. *Global J Pharmacol* 2015; 9(4): 289-295.
- Sivakumar A, Manikandan A, Rajini Raja M and Jayaraman G: *Andrographis paniculata* leaf extracts as potential *N. naja* anti-snake venom. *World J of Pharmacy & Pharmaceutical Sciences* 2015; 4(12): 1036-50.
- Rajeshkumar S, Nagalingam M, Ponnaniakamideen M, Vanaja M and Malarkodi C: Anticancer activity of *Andrographis paniculata* leaves extract against neuroblastoma (Imr-32) and human colon (Ht-29) cancer cell line. *World Journal of Pharmacy and Pharmaceutical Sciences* 2015; 4(6): 1667-1675.
- Madav S, Tripathi HC, Tandan SK and Mishra S: Analgesic, the antipyretic and antiulcerogenic effect of andrographolide. *Ind J Pharm Sci* 1995; 57: 121-125.
- Bu-Chin Y, Chen-Road H, Wang-Chuan C and Juei-Tang C: Antihyperglycemic effect of andrographolide in streptozotocin-induced diabetic rats. *Planta Medica* 2003; 69: 1075-1079.
- Sheeja K and Kuttan G: Activation of cytotoxic T lymphocyte responses and attenuation of tumor growth *in-vivo* by *A. paniculata* extract and andrographolide. *Immunopharmacol Immunotoxicol* 2007; 29: 81-93.
- Calabrese C, Berman SH, Babish JG, MaX, Shinto L, Dorr M, Wells K, Wenner GA and Standish LJ: A phase I trial of andrographolide in HIV positive patients and normal volunteers. *Phytother Res* 2000; 14: 333-338.
- Yoopan N, Thisoda P, Rangkadilok N, Sahasitawat S, Pholphana N, Ruchirawat S and Satayavivad J: Cardiovascular effects of 14-deoxy-11, 12-didehydro andrographolide and *Andrographis paniculata* extracts. *Planta Med* 2007; 73: 503-511.
- Trivedi NP, Rawal UM and Patel BP: Hepatoprotective effect of andrographolide against hexachloro cyclohexane-induced oxidative injury. *Integr Cancer Ther* 2007; 6: 271-280.
- Sharmila N and Gomathi N: Antibacterial, antioxidant activity and phytochemical studies of *Crossandra infundibuliformis* leaf extracts. *International Journal of Phytomedicine* 2011; 3: 151-156.
- Harborne JB: Phytochemical methods. In Chapman and Hall, New York, Edition 2nd, 1984: 21-26.
- Gayathri V and Kiruba D: Preliminary phytochemical analysis of leaf powder extracts of *Psidium guajava* L. *International Journal of Pharmacognosy and Phytochemical Research* 2014; 6(2): 332-334.
- Perez C, Pauli M and Bazerque P: An antibiotic assay by agar-well diffusion method. *Acta Biologica et Medecine Experimental* 1990; 15: 113-115.
- Tavazzi B, Amorini AM, Fazzina G, Di-Pierro D, Tuttobene M and Giardina B: Oxidative stress induces impairment of human erythrocyte energy metabolism through the oxygen radical-mediated direct activation of AMP-deaminase. *Journal of Biology and Chemistry* 2001; 276: 48083-48092.
- Dluya T, Daniel D and Yaduma Gaiuson W: Comparative biochemical evaluation of leaf extracts of *Ficus sycomorus* and *Piliostigma thonningii* plant. *Journal of Medicinal Plants Studies* 2015; 3(5): 32-37.
- Chakraborty D and Shah B: Antimicrobial, antioxidative and antihemolytic activity of *Piper betel* leaf extracts. *Int J Pharm Sci* 2011; 3(3): 192-199.
- Kupier-Goodman T and Scott PM: Risk assessment of mycotoxin ochratoxin A. *Biomedical and Environmental Sciences* 1989; 2: 179-248.
- Nickavar B, Kamalinejad M, Haj-Yahya M and Shafaghi B: Comparison of the free radical scavenging activity of six Iranian *Achillea species*. *Pharmaceutical Biology* 2006; 44: 208-12.

28. James O and Alewo IM: *In-vitro*, antihaemolytic activity of *Gymnema sylvestre* extracts against hydrogen peroxide (H₂O₂) induced haemolysis in human erythrocytes. American Journal of Phytomedicine and Clinical Therapeutics 2014; 2(7): 861-869.
29. Kumarappan C, Jaswanth A and Kumarasunderi K: Antihaemolytic and snake venom are neutralizing effect of

- some Indian medicinal plants. Asian Pacific Journal of Tropical Medicine 2011; 743-747.
30. Sati SC, Takuli P, Kuma P and Khulbe K: Antibacterial activity of three medicinal plants of Kumaun Himalaya against some pathogenic bacteria. International Journal of Pharma Sciences and Research 2015; 6(11): 1361-68.

How to cite this article:

Kadali VN, Ramesh T, Pola SR and Sandeep BV: *In-vitro* anti-haemolytic, anti-bacterial and phytoconstituents investigation of *Andrographis paniculata* methanolic leaf extract. Int J Life Sci & Rev 2016; 2(5): 102-08. doi: 10.13040/IJPSR.0975-8232.IJLSR.2(5).102-08.

All © 2015 are reserved by International Journal of Life Sciences and Review. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **Android OS** based mobile. Scan **QR** Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)