



Received on 12 August 2018; received in revised form, 25 September 2018; accepted, 30 September 2018; published 01 October 2018

## A REVIEW ON SOME MEDICINAL PLANT HAVING ANTIDIABETIC POTENTIAL

Archana Dongray<sup>\*1</sup>, Sarika Nigam<sup>1</sup>, Dilip Kumar Chanchal<sup>2</sup>, Saurabh Chaudhary<sup>2</sup> and Shahbaz Khan<sup>3</sup>

College of Pharmacy<sup>1</sup>, SRGI, Ambabai, Jhansi - 284002, Uttar Pradesh, India.

Department of Pharmacognosy<sup>2</sup>, Institute of Pharmacy, Bundelkhand University, Jhansi - 284128, Uttar Pradesh, India.

Mexcell Pharmaceutical Pvt. Ltd.<sup>3</sup>, Gumnawara, Medical College, Jhansi - 284128, Uttar Pradesh, India.

**ABSTRACT:** Diabetes is a metabolic disorder of the endocrine system. The medicinal plant has used for the treatment of diabetes throughout the world. Several medicinal plants are known to treat diabetes, and they have no side effect. The medicinal plant performed a good clinical practice and is showing a bright future in the treatment of diabetes mellitus. Many studies have confirmed the benefits of the medicinal plant with a hypoglycemic effect in the management of diabetes mellitus. This present review evaluates the medicinal plant used for anti-diabetic activity. The present review profile gives information about the plant's scientific name, common name, family, and the part of the plant used to treat diabetes mellitus.

**Keywords:** Medicinal plant, Diabetic potential

### Correspondence to Author:

**Ms. Archana Dongray**

Assistant Professor, College of Pharmacy, SRGI, Ambabai, Jhansi - 284002, Uttar Pradesh, India.

**E-mail:** archanaadongray@gmail.com

**INTRODUCTION:** Medicinal plant continues to be an important therapeutic aid for alleviating ailments of humankind. Over the last 2500 years, there has been a very strong traditional system of medicine such as Chinese, Ayurvedic, and the Unani, born and practiced more in the eastern continent. These traditions are still flourishing, since; approximately 80% of the people in the developing countries rely on these systems of medicine for their primary health care needs medicinal plants are good sources for new safe, biodegradable and renewable drugs<sup>1</sup>.

**History of Use of Traditional Herbal Medicines:** By definition, 'traditional' use of herbal medicines implies substantial historical use, and this is certainly true for many products that are available as 'traditional herb medicines.' In many developing countries, a large proportion of the population relies on traditional practitioners and their armamentarium of medicinal plants to meet healthcare needs. In Germany, for example, where herbal products are sold as 'phytomedicines,' they are subject to the same criteria for efficacy, safety, and quality as are other drug products<sup>2</sup>.

**The Role of Herbal Medicines in Traditional Healing:** The pharmacological treatment of disease began long ago with the use of herbs<sup>3</sup>. Methods of folk healing throughout the world commonly used herbs as part of their tradition. Some of these traditions are briefly described below, providing some examples of the array of important healing

	<p><b>QUICK RESPONSE CODE</b></p>
	<p><b>DOI:</b> 10.13040/IJPSR.0975-8232.IJLSR.4(10).155-62</p>
<p>The article can be accessed online on <a href="http://www.ijlsr.com">www.ijlsr.com</a></p>	
<p><b>DOI link:</b> <a href="http://dx.doi.org/10.13040/IJPSR.0975-8232.IJLSR.4(10).155-62">http://dx.doi.org/10.13040/IJPSR.0975-8232.IJLSR.4(10).155-62</a></p>	

practices around the world that used herbs for this purpose.

**Indian Traditional Medicine:** Ayurveda is a medical system primarily practiced in India that has been known for nearly 5000 years. It includes diet and herbal remedies while emphasizing the body, mind, and spirit in disease prevention and treatment <sup>4</sup>.

**Japanese Traditional Medicine:** Many herbal remedies found their way from China into the Japanese systems of traditional healing. Herbs native to Japan were classified in the first pharmacopeia of Japanese traditional medicine in the ninth century <sup>5</sup>.

**Traditional Chinese Medicine:** Traditional Chinese medicine has been used by Chinese people from ancient times. Although animal and mineral materials have been used, the primary source of remedies is botanical. Of the more than 12 000 items used by traditional healers, about 500 are in common use. Traditional Chinese medicine is still in common use in China. More than half the population regularly uses traditional remedies, with the highest prevalence of use in rural areas. About 5000 traditional remedies are available in China; they account for approximately one-fifth of the entire Chinese pharmaceutical market <sup>6</sup>.

**History of Herbs in Medicine and Pharmacy:** Herbs were our first source of medicine, and their use predates written history by several thousand years. No one knows when humans first used plants for medicine, but pollens of at least six medicinal plants were found in a Neanderthal burial site estimated to be at least 60,000 <sup>7</sup>. The early history of medicine parallels the history of herbal medicine: the first books written about medicine were also the first books written about herbs, including Chinese texts from 5000 yr ago, such as the famous herbal of the Yellow Emperor and the Egyptian text Ebers papyrus, written 3500 yr ago.

In Western medicine, the father of modern medicine, Theophrastus, is also the father of modern botany. Theophrastus published the first book describing plants in detail in 320 BC, which was also the first Western book about their medicinal uses. Herbal medicine has been at the heart of medicine in every culture in the world and at every time throughout history.

Today, according to the World Health Organization (WHO), more than 80% of the world's population relies on traditional medicines, mostly plant-based, as their main source of health care <sup>8</sup>. This figure includes not only the large populations of China and India and all of the less developed countries of the world but also many modern nations. Even in the United States, approx 25% of our prescription medicines are still extracted from plants or are synthetic copies of plant chemicals <sup>8</sup>, and at least 57% of our top prescription medicines are derived in some way from plants, including semisynthetic, in which plant chemicals are used as building blocks for synthetic drugs <sup>9</sup>.

Diabetes mellitus is a group of metabolic disorder characterized by hyperglycemia resulting from defects in insulin secretion, action, or both. It is made up of two types: Type I and type II. Type I diabetes often referred to as juvenile diabetes, is insulin dependent and known to affect only 5% of the diabetic population. Type II, which is non-insulin dependent, usually develops in adults over the age of 40. It has already been established that chronic hyperglycemia of diabetes is associated with long term damage, dysfunction, and eventually the failure of organs, especially the eyes, kidneys, nerves, heart, and blood vessels <sup>10</sup>. It is estimated that 25% of the world population is affected by this disease. Currently, available therapy for diabetes includes insulin and various oral hypoglycemic agents such as sulfonylureas, metformin, glucosidase inhibitors, troglitazone, *etc.* But these are reported to produce serious adverse side effects such as liver problems, lactic acidosis and diarrhea <sup>11</sup>. It is currently affecting around 143 million people <sup>12</sup>, and the number of those affected is increasing day by day, by 2030 it is predicted to reach 366 million populations worldwide <sup>13</sup>.

According to the World Health Organization (WHO), about 65-80% of the world's population in developing countries depends essentially on plants and plant-derived compounds for their primary healthcare needs. According to the National Medicinal Board, Govt. of India, 17,000 to 18,000 number of species of flowering plants are estimated of which 6,000 to 7,000, species are found to have medicinal usage in folk and documented System of medicine like Ayurveda, Siddha, Unani, and Homeopathy <sup>14</sup>.

**Type of Diabetes:**

- ✓ Type 1 (Insulin dependent diabetes mellitus).
- ✓ Type 2 (Non insulin dependent diabetes mellitus).
- ✓ Type 3 (Gestational diabetes, mellitus).

**Type 1 (Insulin Dependent Diabetes Mellitus):** It is also called as Insulin Dependent Diabetes Mellitus (IDDM). It is due to failure of the body for insulin production (International diabetes federation)<sup>15</sup>.

It is often childhood disease, so it is also called as Juvenile onset diabetes mellitus.

**Insulin Dependent Diabetes Mellitus (or Juvenile Onset Diabetes)**

- ✓ A most common type of diabetes in children
- ✓ An autoimmune disorder in which the immune system attacks and destroys the part of the pancreas that makes insulin.
- ✓ Symptoms include increased thirst, increased urination, increased hunger, weight loss, fatigue,
- ✓ Abdominal pain.
- ✓ Requires daily insulin injections to control blood sugar.
- ✓ Can't be prevented (autoimmune disorder.)

**TABLE 1: ANALYSIS OF REMEDIES OBTAINED FROM DIFFERENT PLANT PARTS FOR DIABETES MELLITUS**

S. no.	Botanical name	Common name	Family	Active chemical constituents	Part used
1	<i>Acacia Arabica</i>	Indian gum	Fabaceae	Polyphenol, Tannin <sup>7-12</sup>	Seed, bark
2	<i>Aegle mameelos</i>	Golden Apple	Rutaceae	Aegeline 2, Coumarin, Flavonoid, Alkaloid <sup>13-15</sup>	Leaf, seed, Fruit
3	<i>Allium cepa</i>	Onion	Liliaceae	Allyl propyl disulfide, S- methyl cysteine sulphoxide <sup>12,16, 17</sup>	Bulb
4	<i>Aloe barbadensis</i>	Barbados Aloe	Asphodelaceae	Lophenol, 24-methyl-lophenol, 24-Ethyllophenol <sup>8, 19</sup>	Leaf
5	<i>Allium sativum</i> L.	Garlic	Alliaceae	Diallyl disulfide oxide, Ajoene, Allyl propyl disulfide, S-allyl cysteine, S-allyl mercaptocysteine <sup>20-22</sup>	Root
6	<i>Azadirachta indica</i>	Neem	Meliaceae	Nimbidine <sup>11, 12, 23</sup>	Leaf, seed
7	<i>A. heterophyllus</i>	Jackfruit	Moraceae	Sapogenin <sup>11, 12, 24, 25</sup>	Sapogenin
8	<i>Aloe vera</i>	Barbados aloe	Liliaceae	Pseudoprotinosaponin, Prototinosaponin <sup>18, 25, 26</sup>	Leaf
9	<i>Annona Sqamosa</i>		Annonaceae <sup>26-29</sup>		fruit peel
10	<i>A. sphaerocephala</i>	Wormwood	Asteraceae	Polysaccharide <sup>11, 12, 30, 31</sup>	Fruit
11	<i>Abelmoschus esculentus</i>	Gumbo	Malvaceae	Carbohydrate, Gum, Mucilage, Protein, Phytosterol, Flavonoid, Tannin, Phenolics, Volatile oil <sup>11</sup>	Fruit
12	<i>Abrus precatorius</i> L.	Jequirity	Fabaceae <sup>32, 33</sup>		Seed
13	<i>Abutilon crispum</i> (L.)	Nela benda	Malvaceae <sup>34</sup>		Leaf
14	<i>Butea monosperma</i>	Bastard teak	Fabaceae	Butein, Palasonin, Stigmasterol-3 β-D-glucopyranoside <sup>11</sup>	Fruit
15	<i>Beta vulgaris</i>	Beetroot	Chenopodiaceae	Sugar beet pectin, Polydextrose <sup>12, 35</sup>	Whole plant
16	<i>Biophytu. Sensitivum</i>	Sikerpud	Oxalidaceae <sup>12, 26, 36</sup>	-	Whole plant
17	<i>Brassica juncea</i>	Mustard	Brassicaceae	Isorhamnetin diglucoside <sup>37</sup>	Seed, Leaf
18	<i>Brassica nigra</i> L	Black mustard	Cruciferae <sup>38</sup>		Whole plant
19	<i>Bauhinia acuminata</i>	White kachnar	Caesalpiniaceae <sup>39</sup>		Leaf
20	<i>Boerhaavia diffusa</i> L.	Tar vine	Nyctaginaceae <sup>38</sup>		Leaf
21	<i>Bridelia ferruginea</i>	Balli chettu	Phyllanthaceae <sup>40</sup>		Leaf, fruit, stem bark, roots
22	<i>Carica papaya</i>	Papaya	Caricaceae	Saponin, Tannin, Alkaloid, Flavonoid <sup>11</sup>	Fruit
23	<i>Coccinia</i>	Ivy-gourd	Cucurbitaceae	B-amyrin, Lupeol, Cucurbitacin B <sup>11, 12,</sup>	Fruit

	<i>indica</i>				
24	<i>Curcuma longa</i>	Turmeric	Zingiberaceae	Curcuminoid <sup>12</sup>	Root
25	<i>Cuminum cyminum</i>	Cumin seed	Apiaceae	Aldehyde <sup>12, 42</sup>	Seed
26	<i>C. sativum</i>	Coriander	Apiaceae	Alanine <sup>12, 42</sup>	Leaf
27	<i>C. zeylanicum</i>	Cinnamon	Lauraceae	Cinnamaldehyde <sup>11, 12</sup>	Leaf, Bark
28	<i>Catharanthus</i>	Red periwinkle	Apocynaceae	Vinculin, Alkaloid <sup>12</sup>	Whole plant
	<i>Roseus</i>				
29	<i>Capsicum</i>	Chilli	Solanaceae	Capsaicin <sup>43</sup>	Fruit
	<i>frutescens</i>				
30	<i>Diospyros lotus</i>	Date plum	Ebenaceae	Phenolics <sup>11</sup>	Fruit
31	<i>Datura stramonium</i>	Thorn apple	Solanaceae <sup>33</sup>		Seed
32	<i>Emblica officinalis</i>	Amla	Euphorbiaceae	Tannoid <sup>11, 12</sup>	Fruit
33	<i>Eucalyptus</i>	Blue gum	Myrtaceae	Calytoside <sup>11, 12, 44</sup>	Leaf
	<i>globules</i>				
34	<i>Erythrina variegata</i>	Sunshine tree	Fabaceae		Root
35	<i>Ficus bengalensis</i>	Banyan tree	Moraceae	Leucopelargonidin <sup>11, 12, 46</sup>	Bark
36	<i>Ficus carica</i>	Anjir	Moraceae	Invert sugar <sup>11, 12, 46</sup>	Leaf, fruit
37	<i>Gymnema</i>	Sugar	Asclepiadaceae	Gymnemic acid, Gymnema, Saponin <sup>47, 12</sup>	Leaf
	<i>sylvestre</i>	Destroyer			
38	<i>Grifola frondosa</i>	Maitake	Meripilaceae	Disaccharide <sup>11</sup>	Fruit
39	<i>Glycine max</i>	Soya beans	Fabaceae	3-O-methyl-D-chiro-inositol <sup>9-12</sup>	Seed
40	<i>H. auriculata</i>	Talmakhana	Acanthaceae <sup>12</sup>	-	Whole plant
41	<i>Helicteres</i>	East Indian	Sterculiaceae	Steroid, Terpenoid, Alkaloid, Carbohydrate, Phenolics <sup>11</sup>	Fruit
	<i>isora</i>	screw tree			
42	<i>Ibervillea</i>	Huereque	Cucurbitaceae	Monoglyceride (MG), Fatty acid <sup>41, 11, 12</sup>	Root
	<i>sonorae</i>				
43	<i>Ipomoea batatas</i>	Sweet potato	Convolvulaceae <sup>48</sup>		Leaf
44	<i>Justicia glauca</i>	Water-willow	Acanthaceae <sup>49</sup>		Plant
45	<i>Jatropha curcas</i>	Barbados nut	Euphorbiaceae	Tannin <sup>11, 12, 49</sup>	Whole plant
46	<i>Lyophyllum</i>	<i>Lyophyllum</i>	Lyophyllaceae	Polysaccharide <sup>11</sup>	Fruit
	<i>decastes</i>	<i>Decastes</i>			
47	<i>L. sechellarum</i>	Sea coconut	Palmae	Carbohydrate <sup>11</sup>	Fruit
48	<i>Lycium barbarum</i>	Chirchita	Solanaceae	Polysaccharide <sup>50, 51</sup>	Fruit
49	<i>Limonia acidissima</i>	Wood apple	Rutaceae	Polysaccharide <sup>11, 12</sup>	Fruit
50	<i>Momordica</i>	Bitter melon	Cucurbitaceae	Charantin, Momordicin, Galactose binding, Lectin, Non-bitter, Diosgenin, Cholesterol, lanosterol, $\beta$ -sitosterol, Cucurbitacin glycoside <sup>52, 53, 11, 12</sup>	Whole plant
	<i>charantia</i>				
51	<i>Mentha</i>	Peppermint	Lamiaceae	Essential oil, Terpen, Flavonoid. Vanadium, Zinc, Chromium, Copper, Iron, Potassium, Sodium, Nickel <sup>54, 12</sup>	Leaf
	<i>piperita</i>				
52	<i>Mangifera</i>	Mango	Anacardiaceae	Mangiferin, Phenolics, Flavonoid <sup>55, 11, 12</sup>	Leaf, stem Bark, fruit
	<i>indica</i>	tree			
53	<i>Musa paradisiaca</i>	Banana	Musaceae	Dietary fibre, Pectin <sup>11, 12</sup>	Fruit
54	<i>Musa sapientum</i>	Sweet banana	Musaceae	Flavonoid, Steroid, Glycoside <sup>56</sup>	Flower
55	<i>Nigella sativa</i>	Roman	Ranunculaceae	Thymoquinone <sup>57, 12</sup>	Whole plant
	<i>Coriander</i>				
56	<i>Opuntia dillenii</i>	Prickly pear	Cactaceae	Polysaccharide <sup>12</sup>	Fruit
57	<i>Ocimum</i>	Holy basil	Lamiaceae	Eugenol (1-hydroxy-2-methoxy-4- 26] Allylbenzene <sup>58, 12</sup>	Leaf
	<i>sanctum</i>				
58	<i>Physalis</i>	Strawberry	Solanaceae	Polysaccharide <sup>11, 12</sup>	
	<i>alkekengi</i>	Tomato			
59	<i>Persea americana</i>	Avocado	Lauraceae	Fat, Protein, Vitamin, Mineral <sup>11, 12</sup>	Fruit
60	<i>Psidium guajava</i>	Guava	Myrtaceae	Terpen, Flavonoid, Strictinin,	Leaf,

				Isostrictinin, Pedunculagin, Polysaccharide <sup>59</sup> Tannin <sup>11, 12</sup>	Fruit
61	<i>Phyllanthus emblica</i> ; <i>P. Acidus</i>	Indian gooseberry	Euphorbiaceae		Fruit
62	<i>Piper betle</i>	Pan	Piperaceae <sup>60</sup>		Leaf
63	<i>Piper longum</i>		Piperaceae <sup>11, 60, 12</sup>		Root
64	<i>Punica granatum</i>	Pomegranate	Punicaceae	Tannin <sup>12</sup>	Fruit
65	<i>Panax ginseng</i>	Ginseng	Araliaceae	Saponin <sup>12</sup>	Fruit
66	<i>Rhus coriaria</i>	Sicilian Sumac	Anacardiaceae	Limonene, Nonanal, Dec-2 (Z)-enal <sup>11, 12</sup>	Fruit
67	<i>Thespesia populnea</i>	Portia tree	Malvaceae	Populnetin, Herbacetin, Populneol, Quercetin <sup>12</sup>	Fruit
68	<i>Terminalia chebula</i>	Chebolic myrobalan	Combretaceae	Shikimic, Gallic, Triacontanoic, Palmitic acid, $\beta$ -sitosterol, Daucosterol <sup>61, 12</sup>	Stem bark
69	<i>Terminalia Catappa</i>	Indian almond	Combretaceae	Petroleum ether, Methanolic, Aqueous <sup>11, 12, 61</sup>	Fruit
70	<i>Turnera diffusa</i>	Damiana	Turneraceae	Flavonoid, Terpen <sup>12</sup>	Leaf
71	<i>Tamarindus indica</i>	Tatul tree	Fabaceae	Flavonoid, Polysaccharide <sup>10, 11, 12</sup>	Seed, Fruit
72	<i>Triticum vulgare</i>	Wheat	Poaceae	Albumin <sup>11,12</sup>	Whole plant
73	<i>V. angustifolium</i>	Wild blueberry	Ericaceae	Phenolic <sup>10, 11, 12</sup>	Fruit
74	<i>Viburnum opulus</i>	Cranberry bush	Caprifoliaceae	Tannin <sup>11, 12</sup>	Fruit
75	<i>V. myrtillus</i>	Bilberry	Ericaceae	Anthocyanoside <sup>10, 11, 12</sup>	Leaf, Fruit
76	<i>W. somnifera</i>	Winter cherry	Solanaceae	Withanolide, Alkaloid <sup>62, 10, 11</sup>	Leaf
77	<i>Withania coagulans</i>	Vegetable Rennet	Solanaceae	Milk-coagulating enzyme, Esterase, Fatty oil, Essential oil, Alkaloid <sup>62, 10, 11</sup>	Fruit
78	<i>Xanthocercis zambesiaca</i>	Nyala tree	Fabaceae	Fagomine, 4-O-beta-Dglucopyranosylfagomine, Castanospermine <sup>7, 9, 10</sup>	Leaf
79	<i>Zingiber officinale</i>	Ginger	Zingiberaceae	Gingerol, Ethanol <sup>63, 12</sup>	Bulb
80	<i>Ziziphus spinachristi</i>	Christ thorn	Rhamnaceae	Christinin-A, Fatty acid <sup>64, 65</sup>	Leaf

**DISCUSSION:** Diabetes mellitus is spreading alarmingly throughout the world and three fourths of the world populations and considered as a major cause of high economic loss which can, in turn, impede the development of nations. Moreover, uncontrolled diabetes leads to many chronic complications such as blindness, heart disease, and renal failure, etc. Diabetes is increasing day – by – day, presently, insulin is the only drug before ailing patients. However, some crude drug of herbal origin is in use of the market. The promising crude drug must be analyzed in clinically manifested hyperglycemia in the wake of a thorough investigation of ethnomedicinal anti-diabetic herbs.

The study revealed that 80 plant species belonging to 49 families were generally used for the treatment of diabetes. The majority of the experiments confirmed the benefits of medicinal plants with hypoglycemic effects in the management of diabetes mellitus. Among the plants used for diabetes, *Annona squamosa*, *Momordica charantia*,

*Egyptian Morus alba*, *Lycium barbarum*, *Allium sativum*, and *Aegle marmelose* seems to be most common plants used to treat diabetes and are available everywhere. The most commonly involved active constituents are flavonoid, tannin, phenolics, and alkaloid. Numerous mechanisms of actions have been proposed for these plant extracts. Some hypotheses relate to their effects on the activity of pancreatic  $\beta$ -cells (synthesis, release) or the increase of the insulin sensitivity or the insulin-like activity of the plant extracts. All of these actions may be responsible for the reduction or abolition of diabetic complications.

**CONCLUSION:** Long before the use of insulin became common; indigenous remedies were used for the treatment of diabetes mellitus. There has been an increasing demand from patients for the use of natural products with antidiabetic activity. This is largely because insulin cannot be used orally and insulin injections are associated with the risk of hypoglycemia and impairment of hepatic

and other body functions. The undesirable side effects and contraindications of synthetic drugs and the fact that they are not suitable for the use during pregnancy have made scientists look towards hypoglycaemic agents of plant origin.

The present review has presented comprehensive details of anti-diabetic plants used in the treatment of diabetes mellitus. Some of these plant-derived medicines, however, offer the potential for cost-effective management of diabetes through dietary interventions, nutrient supplementation, and combination therapies with synthetic drugs in the short term, and as the sole medication from natural sources over the long term. The presences of bioactive chemicals are mainly responsible for this anti-diabetic action.

However, many other active agents obtained from plants have not been well characterized. More investigations must be carried out to evaluate the mechanism of action of medicinal plants with an anti-diabetic effect.

**ACKNOWLEDGEMENT:** The authors thankful with our deepest core of heart to Mr. Dilip Kumar Chanchal for his valuable guidance.

**CONFLICT OF INTEREST:** Nil

## REFERENCES:

1. Tsay HS and Agrawal DC: Tissue Culture Technology of Chinese Medicinal Plant Resources in Taiwan and their Sustainable Utilization. *Int J App Sci Eng* 2005; 3:215-223.
2. Shaw D: Risks or remedies? Safety aspects of herbal remedies. *J Roy Soc Med* 1998; 91: 294-296.
3. Schulz V, Hänsel R and Tyler VE: Rational Phytotherapy. A Physician's Guide to Herbal Medicine, Berlin, Springer-Verlag, Edition 4<sup>th</sup>, 2001.
4. Morgan K: Medicine of the Gods: Basic Principles of Ayurvedic Medicine 2002 [<http://www.compulink.co.uk/~mandrake/ayurveda.htm>]
5. Saito H: Regulation of herbal medicines in Japan. *Pharmacol. Regul.* 2000; 41: 515-519.
6. Li L: Opportunity and challenge of traditional Chinese medicine in the face of the entrance to WTO (World Trade Organization)]. *Chin Inform Trad Chin Med* 2000; 7: 7-8 (in Chinese).
7. Soleck RSV and Shanidar IV: A Neanderthal flower burial in Northern Iraq. *Science*.
8. Farnsworth NR, Akerele O and Bingel AS: Medicinal plants in therapy. *Bull World Health Org* 1985; 63: 965-981.
9. Grifo F, Newman D and Fairfield AS: The origins of prescription drugs. In: Grifo F, Rosenthal J, eds. *Biodiversity and Human Health*. Island Press, Washington, DC, 1997: 131-163.

10. Huang THW, Peng G, Kota BP, Li GQ, Yamahara J and Roufogalis BD: Anti-diabetic action of *Punica granatum* flower extract: activation of PPAR-c and identification of an active component. *Toxicol App Pharmacol* 2005; 207: 160-169.
11. Rajalakshmi M, Eliza J, Priya CE, Nirmala A and Daisy P: Antidiabetic properties of *Tinospora cordifolia* stem extracts on streptozotocin-induced diabetic rats. *Afr J Pharm Pharmacol* 2009; 3(5): 171-180.
12. Ponnusamy S, Ravindran R, Zinjard S, Bhargava S and Kumar AR: Evaluation of traditional Indian antidiabetic medicinal plants for human pancreatic amylase inhibitory effect *in-vitro*. *Evid Based Complementary Altern Med* 2011; 1-10.
13. Swargiary A, Boro H, Bramha BK and Rahman S: Ethnobotanical study of the anti-diabetic medicinal plant used by the local people of Kokrajhar District of Bodoland Territorial Council, India *Journal of Medicinal plant Studies* Year-2013; 1(5).
14. Morgan K: Medicine of the Gods: Basic Principles of Ayurvedic Medicine 2002 [<http://www.compulink.co.uk/~mandrake/ayurveda.htm>]
15. Nojima H, Kimura I, Chen FJ, Sugihara Y, Haruno MV and Kato A: Antihyperglycemic effects of N containing sugars from *Xanthocercis zambesiaca*, *Morus bombycis*, *Aglaonema treubii*, and *Castanospermum australe* in streptozotocin-diabetic mice. *J Nat Prod* 1998; 61(3): 397-400.
16. Hatapakki BC, Suresh HM, Bhoomannavar V and Shivkumar SI: Effect of *Cassia auriculata* Linn. Flowers against alloxan-induced diabetes in rats. *J Nat Remedies* 2005; 5(2): 132-136.
17. Maghrani M, Michel JB and Eddouks M: Hypoglycaemic activity of *Retama raetam* in rats. *Phytotherapy Res* 2005; 19: 125-128.
18. Kang MJ, Kim JI, Yoon SY, Kim JC and Cha IJ: Pinitol from soybeans reduces postprandial blood glucose in patients with type 2 diabetes mellitus. *J Med Food* 2006; 9(2): 182-186.
19. Vikrant A and Sharma R: A review on fruits having anti-diabetic potential. *Journal of Chemical and Pharmaceutical Research*. *J Chem Pharm Res* 2011; 3(2): 204-212.
20. Makheswari MU and Sudarsanam D: Database on Anti-diabetic indigenous plants of Tamil Nadu, India. *Int J Pharma Sci Res* 2012; 3(2): 287-293.
21. Kamalakkannan N and Prince PS: The effect of *Aegle marmelos* fruit extract in streptozotocin diabetes: a histopathological study. *J Herbal Pharmacother* 2005; 5: 87-96.
22. Kesari AN, Gupta RK, Singh SK, Diwakar S and Watal G: Hypo-glycemic and antihyperglycemic activity of *Aegle marmelos* seed extract in normal and diabetic rats. *J Ethnopharmacol* 2006; 107: 374-379.
23. Narendar T, Shweta S, Tiwari P, Papi Reddy K, Khaliq T, Prathipati P: Antihyperglycemic and an antidyslipidemic agent from *Aegle marmelos*. *Bioorg Med Chem Lett* 2007; 17(6): 1808-1811.
24. Fetrow CW and Avila JR: Professional's Handbook of Complementary and Alternative Medicines. Springhouse, PA: Springhouse Corporation, 1999.
25. Hattori A, Yamada N, Nishikawa T, Fukuda H and Fujino T: Antidiabetic effects of ajoene in genetically diabetic KK-A(y) mice. *J Nutr Sci Vitamil* 2005; 51(5): 382-384.
26. Pillai VR and Santhakumari G: Hypoglycaemic activity of *Melia azadirachta* Linn. (Neem). *Indian J Med Res* 1981; 74: 931.

27. Tanaka M, Misawa E, Ito Y, Habara N, Nomaguchi K and Yamada M: Identification of five phytosterols from Aloe vera gel as anti-diabetic compounds. *Biol Pharm Bull* 2006; 29: 1418-1422.
28. Liu CT, Wong PL, Lii CK, Hse H and Sheen LY: Antidiabetic effect of garlic oil but not diallyl disulfide in rats with streptozotocin-induced diabetes. *Food Chem Toxicol* 2006; 44: 1377-1384.
29. El-Demerdash FM, Yousef MI, El-Naga NI. Biochemical study on the hypoglycemic effects of onion and garlic in alloxan-induced diabetic rats. *Food Chem Toxicol* 2005; 43: 57-63.
30. Eidi A, Eidi M and Esmaeili E: Antidiabetic effect of garlic (*Allium sativum* L.) in normal and streptozotocin induced diabetic rats. *Phytomedicine* 2006; 13: 624-629.
31. Schwab U, Louheranta A, Törrönen A and Uusitupa M: Impact of sugar beet pectin and polydextrose on fasting and postprandial glycemia and fasting concentrations of serum total and lipoprotein lipids in middle-aged subjects with abnormal glucose metabolism. *Eur J Clin Nutr* 2006; 60(9): 1073-1080.
32. Bnouham M, Ziyat A, Mekhfi H, Tahri A and Legssyer A: Medicinal plants with potential anti-diabetic activity-a review of ten years of herbal medicine research (1990-2000). *Int J Diabetes Metab* 2006; 14: 125.
33. Ayodhya S, Kusum S and Anjali S: Hypoglycaemic activity of different extracts of various herbal plants Singh. *Int J Ayurveda Res Pharm* 2010; 1(1): 212-224.
34. Fröde TS and Medeiros YS: Animal models to test drugs with potential anti-diabetic activity. *J Ethnopharmacol* 2008; 115: 173-183.
35. Gupta RK, Kesari AN, Watal G, Murthy PS, Chandra R and Tandon V: Nutritional and hypoglycemic effect of fruit pulp of *Annona squamosa* in normal healthy and alloxan-induced diabetic rabbits. *Annals Nut Metabol* 2005b; 49: 407-413.
36. Kaleem M, Asif M, Ahmed QU and Bano B: Antidiabetic and antioxidant activity of *Annona squamosa* extract in streptozotocin-induced diabetic rats. *Singapore Med J* 2006; 47: 670-675.
37. Ojewole JA: Antinociceptive, anti-inflammatory and anti-diabetic effects of *Bryophyllum pinnatum* (Crassulaceae) aqueous leaf extract. *J Ethnopharmacol* 2005a; 99: 13-19.
38. Singh LW: Traditional medicinal plants of Manipur as antidiabetics. *J Med Plant Res* 2011; 5(5): 677-687.
39. Shirwaikar A, Rajendran K and Punitha IS: Antidiabetic activity of alcoholic stem extract of *Coscinium fenestratum* in streptozotocin-nicotinamide-induced type 2 diabetic rats. *J Ethnopharmacol* 2005; 97: 369-374.
40. Bhatia M, Siddiquina B and Gupta S: *A. Precatorius* (L): An evaluation of traditional herb. *Indo American Journal of Pharmaceutics Research AJPR* 2013; 4: 3295-3315.
41. Verma BV and ANV: Plant with antidiabetic activities and their medicinal values of review article. *International Research Journal of Pharmacy* 2012; 3(3).
42. Padal SB, Murty PP, Rao DS and Venkaiah M: Ethnomedicinal plant from Paderu of the division of Visakhapatnam district, A.P, Indian Journal of Phytology 2010; 2(8): 70-91.
43. Tan BK, Tan CH and Pushparaj PN: Anti-diabetic activity of the semi-purified fractions of *Averrhoa bilimbi* in high-fat diet fed-streptozotocin-induced diabetic rats. *J Life Sci* 2005; 76: 2827-2839.
44. Kubo H, Kobayashi J, Higashiyama K, Kamei J, Fujii Y and Ohmiya S: The hypoglycemic effect of (7R\*, 9aS\*)-7-phenyl-octahydroquinolizin-2-one in mice. *Biol Pharm Bull* 2000; 23(9): 1114-1117.
45. Yokozawa T, Kim HY, Cho EJ, Choi JS and Chung HY: Antioxidant effects of isorhamnetin 3, 7-di-O-beta-D-glucopyranoside isolated from mustard leaf (*Brassica juncea*) in rats with streptozotocin-induced diabetes. *Agric Food Chem* 2002; 50(19): 5490-5495.
46. Patel DK, Kumar R, Laloo D and Hemalatha S: Diabetes Mellitus: An Overview on its pharmacological aspects and reported medicinal plants having anti-diabetic activity. *Science Direct Asian Pacific Journal of Tropical Biomedicine* 2012; 411-420.
47. Nguemeta TA, Brusotti G, Caccialanza G and Finzi PV: The genus *Bridelia*: A phytochemical and ethno-pharmacological review. *Journal of Ethnopharmacological* 2009; 124: 339-349.
48. Saxena A and Vikram NK: Role of selected Indian plants in management of type 2 diabetes: a review. *J Altern Complement Med* 2004; 10(2): 369-378.
49. Honda S, Aoki F, Tanaka H, Kishida H, Nishiyama T and Okada SM: Effects of ingested turmeric oleoresin on glucose and lipid metabolisms in obese diabetic mice: a DNA microarray study. *J Agric Food Chem* 2006; 54(24): 9055-9062.
50. Tolan I, Ragoobirsingh D and Morrison EY: Isolation and purification of the hypoglycaemic principle present in *Capsicum frutescens*. *Phytother Res* 2004; 18(1): 95-96.
51. Ali M and Chaudhary N: *Ficus hispida* Linn. A review of its pharmacognostic and ethnomedicinal properties. *Pharmacognosy reviews* 2011; 5(9): 96-102.
52. Bnouham M, Ziyat A, Mekhfi H, Tahri A and Legssyer A: Medicinal plants with a potential anti-diabetic activity-a review of ten years of herbal medicine research (1990-2000). *Int J Diabetes Metab* 2006; 14: 1-25.
53. Sugihara Y, Nojima H, Matsuda H, Murakami T, Yoshikawa M and Kimura I: Antihyperglycemic effects of gymnemic acid IV, a compound derived from *Gymnema sylvestris* leaves in streptozotocin-diabetic mice. *J Asian Nat Prod Res* 2000; 2(4): 321-327.
54. Rawat M and Parmar N: Medicinal plant with antidiabetic potential-A Review. *American-Eurasian J Agric & Environ Sci* 2013; 13(1): 81-94.
55. Muruganandan S, Srinivasan K, Gupta S, Gupta PK and Lal J: Effect of mangiferin on hyperglycemia and atherogenicity in streptozotocin diabetic rats. *J Ethnopharmacol* 2005; 97(3): 497-501.
56. Zhao R, Li Qv and Xiao B: Effect of *Lycium barbarum* polysaccharide on the improvement of insulin resistance in NIDDM rats. *Yakugaku Zasshi J Pharm Society Japan* 2005; 125: 981-988.
57. Wu H, Guo H and Zhao R: Effect of *Lycium barbarum* polysaccharide on the improvement of antioxidant ability and DNA damage in NIDDM rats. *Yakugaku Zasshi J Pharm Society Japan* 2006; 126: 365-3671
58. Shetty AK, Kumar GS, Sambaiah K and Salimath PV. Effect of bitter melon (*Momordica charantia*) on glycaemic status in streptozotocin-induced diabetic rats. *Plant Foods Human Nut* 2005; 60: 109-112.
59. Sathishsekar D and Subramanian S: Beneficial effects of *Momordica charantia* seeds in the treatment of STZ induced diabetes in experimental rats. *Biol Pharm Bull* 2005; 28: 978-983.
60. Dhanabal SP, Sureshkumar M, Ramanathan M and Suresh B: Hypoglycemic effect of ethanolic extract of *Musa sapientum* on alloxan induced diabetes mellitus in rats and its relation with antioxidant potential. *J Herb Pharmacother* 2005; 5(2): 7-19.
61. Ojewole JA: Anti-inflammatory, analgesic and hypoglycemic effects of *Mangifera indica* Linn.

- (Anacardiaceae) stem-bark aqueous extract. Methods Findings Experiment Clin Pharmacol 2005c; 27: 547-554.
62. Dhanabal SP, Sureshkumar M, Ramanathan M and Suresh B: Hypoglycemic effect of ethanolic extract of *Musa sapientum* on alloxan induced diabetes mellitus in rats and its relation with antioxidant potential. J Herb Pharmacother 2005; 5(2): 7-19.
  63. Kanter M: Effects of *Nigella sativa* and its major constituent, thymoquinone on sciatic nerves in experimental diabetic neuropathy. Neurochem Res 2008; 33(1): 87-96.
  64. Prakash P and Gupta N: Therapeutic uses of *Ocimum sanctum* Linn. (Tulsi) with a note on eugenol and its pharmacological actions: a short review. Indian J Physiol Pharmacol 2005; 49: 125-131.
  65. Ojewole JA: Hypoglycemic and hypotensive effects of *P. guajava* Linn. (Myrtaceae) leaf aqueous extract. Methods Findings Experiment Clin Pharmacol 2005d; 27: 689-695.
  66. Santhakumari P, Prakasam A and Pugalendi KV: Anti-hyperglycemic activity of *Piper betle* leaf on streptozotocin-induced diabetic rats. J Med Food 2006; 9: 108-112.
  67. Rao NK and Nammi S: Antidiabetic and renoprotective effects of the chloroform extract of *Terminalia chebula* Retz seeds in streptozotocin-induced diabetic rats. BMC Complement Alternat Med 2006; 6: 17.
  68. Babu PS, Prabuseenivasan S and Ignacimuthu S: Cinnamaldehyde-a potential anti-diabetic agent. ~ 158 ~Phytomed 2007; 14(1): 15-22.
  69. Kato A, Higuchi Y, Goto H, Kizu H, Okamoto T and Asano N: Inhibitory effects of *Zingiber officinale* Roscoe derived components on aldose reductase activity *in-vitro* and *in-vivo*. J Agric Food Chem 2006; 54(18): 6640- 44.
  70. Abdel-Zaher AO, Salim SY, Assaf MH and Abdel-Hady RH: Antidiabetic activity and toxicity of *Zizyphus spinachristi* leaves. J Ethnopharmacol 2005; 101: 129-138.
  71. Mohamed B, Ziyat A, Mekhfi H, Tahri A and Legssyer A: Medicinal plants with potential antidiabetic activity - A review of ten years of herbal medicine research (1990-2000). Int J Diabetes & Metabol 2006; 14: 1-25.

**How to cite this article:**

Dongray A, Nigam S, Chanchal DK, Chaudhary S and Khan S: A review on some medicinal plant having anti-diabetic potential. Int J Life Sci & Rev 2018; 4(10): 155-62. doi: 10.13040/IJPSR.0975-8232.IJLSR.4(10).155-62.

All © 2015 are reserved by International Journal of Life Sciences and Review. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)