

Received on 21 May 2014; received in revised form, 28 July 2014; accepted, 04 December 2014; published 01 January 2015

SYNTHESIS OF SOME NOVEL 2-SUBSTITUTED IMINO-7-SUBSTITUTED PHENYL-3, 7-DIHYDRO-2H-[1, 3] THIAZOLO [5, 4-E] - THIAZIN-5-AMINE AS ANTIFUNGAL ACTIVITY

B. Chaubey¹, Vineet K. Singh^{* 2}, D. Singh¹, Rajeev K. Verma² and Vimal K. Yadav²

Synthetical Organic Research Lab, Department of Chemistry, T.D.P.G. College¹, Jaunpur - 222002, Uttar Pradesh, India.

Prasad Institute of Technology², Jaunpur - 222002, Uttar Pradesh, India.

ABSTRACT: The condensation of 2-imino-5 substituted benzylidene-4-thiazolidinone (1) with thiourea (2) in the presence of glacial acetic acid to gives 2-substituted imino-7-substituted phenyl-3,7-dihydro-2H-[1,3] thiazolo [5,4-e]-thiazin-5-amine (3). The antifungal activities of the compounds (3a-h) were tested against *A.niger* and *A. flavus*. The growth of both fungi *A. niger* and *A. flavus* are inhibited to some extent by all the synthesized compounds. Hence all are antifungal agents.

Keywords: Thiazole, Thiazole derivatives, Substituted Thiazine derivatives, Antifungal activity

Correspondence to Author:


Vineet K. Singh

Prasad Institute of Technology, Jaunpur - 222002, Uttar Pradesh, India.

E-mail: mailme2805@gmail.com

INTRODUCTION: Heterocyclic compounds comprise the major family of organic compounds. These are enormously essential with a wide range of synthetic, pharmaceutical, and industrial applications and are famous for their biological activities. There is an extensive spectrum of biological activities shown by many compounds containing five-membered heterocyclic rings in their structure. The high therapeutic properties of these heterocycles have encouraged the medicinal chemists to synthesize a large number of novel chemotherapeutic agents. These heterocyclic compounds have broadened the scope in remedying various dispositions in clinical medicines. Thiazoles have been reported to show pharmacological activities¹.

Thiazole (1) is a well known heterocyclic compound with two heteroatoms, sulfur, and nitrogen at position 1 and 3. The tetrahydro derivatives of thiazole are known as thiazolidine (2), and oxo-derivative of thiazolidine is called thiazolidinone (3). Thiazolidinone is an important compound of this group.



Thiazoles represent an important class of heterocyclic compounds. The thiazole derivatives are widely used in the field of medicine, industry and agriculture. The derivatives of thiazole have been reported as fungicides²⁻³, Pesticides⁴, insecticides⁵, antibacterial⁶⁻⁷, anti-inflammatory⁸, anticonvulsant⁹, antitubercular¹⁰, antiviral¹¹, and anticancer¹². An attempt is to synthesize some novel 2-substituted imino-7-substituted phenyl - 3, 7-dihydro-2H-[1, 3] thiazolo [5, 4-e] - thiazin-5-

QUICK RESPONSE CODE 	DOI: 10.13040/IJPSR.0975-8232.IJLSR.1(1).20-22
	The article can be accessed online on www.ijlsr.com
DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.IJLSR.1(1).20-22	

amine. The synthesized compounds (3) have been screened for antifungal activity against *A. niger* and *A. flavus*.

MATERIALS AND METHODS: Melting points were taken in open capillary tubes and are uncorrected. The IR spectra were recorded in KBr on Perkin-Elmer-720 spectrophotometer. The ¹H NMR spectra were recorded in CDCl₃ on Varian A-60 D. spectrophotometer. The chemical shifts are recorded in ppm downfield from TMS, which are used as an internal standard.

2-substituted imino-7-substituted phenyl-3, 7-dihydro-2H-[1, 3] thiazolo [5, 4-e] - thiazin-5-amine (3a-h) (General Method).

A 0.05 mol of 2-substituted imino-5-benzylidene-4-thiazolidinone and 0.05 mol thiourea was dissolved in 100 ml of glacial acetic acid by refluxing for 5 hours, it was allowed to cool and the solid obtained was filtered. The crude products were pressed firmly and recrystallized from ethanol: water. The physical data of all the synthesized compounds are given in **Table 1**.

TABLE 1: PHYSICAL DATA OF ALL THE SYNTHESIZED COMPOUNDS

Comp. no	R	X	M.P.(^o C)	Yield (%)	Molecular formula	Elemental Analysis			IR cm^{-1}	¹ H NMR (CDCl ₃)
						C	H	N		
3a	H	H	189	65	C ₁₁ H ₁₀ N ₄ S ₂	50.38	3.81	21.37	3390(NH), 760 (C-S), 1680 (C=N), 3080 (Ar-CH)	7.19-8.25 (m,5H,Ar-H), 5.71 (s,4H,NH),2.81 (s,1H,CH)
3b	H	p-OMe	177	62	C ₁₂ H ₁₂ N ₄ S ₂ O	49.31	4.10	19.18	3380 (NH), 780 (C-S), 1673 (C=N), 3075 (Ar-CH), 1250 (C-OCH ₃)	7.20-8.24 (m,4H,Ar-H), 5.461 (s,4H,NH), 2.69 (s,1H,CH), 3.53 (s,3H,OCH ₃)
3c	H	p-N(Me) ₂	162	64	C ₁₃ H ₁₅ N ₅ S ₂	51.14	4.91	22.95	3383 (NH), 778 (C-S), 3079 (Ar-CH)	7.22-8.26 (m,4H,Ar-H), 5.61 (s,4H,NH), 2.7 (s,1H,CH), 3.79 (s,6H,NMe ₂)
3d	H	p-NO ₂	159	73	C ₁₁ H ₉ N ₅ S ₂ O ₂	42.99	2.93	22.80	3370 (NH), 775 (C-S), 3101 (Ar-CH), 1470 (C-NO ₂)	7.26-8.35 (m,4H,Ar-H),5.48 (s,4H,NH),2.75(s,1H,CH)
3e	Ph	H	197	72	C ₁₇ H ₁₄ N ₄ S ₂	60.35	4.14	16.56	3365 (NH), 770 (C-S), 3098 (Ar-CH)	7.23-8.36 (m,16H,Ar-H),5.52 (s,3H,NH),2.68(s,1H,CH)
3f	Ph	p-OMe	186	66	C ₁₈ H ₁₆ N ₄ S ₂ O	58.69	4.34	15.21	3349 (NH), 772 (C-S), 3083 (Ar-CH), 1265 (C-OCH ₃)	7.25-8.37 (m,9H,Ar-H),5.56 (s,3H,NH),2.73(s,1H,CH), 3.53 (s,3H,OCH ₃)
3g	Ph	p-N(Me) ₂	178	64	C ₁₉ H ₁₉ N ₅ S ₂	59.84	4.98	18.37	3384 (NH), 762 (C-S), 3075 (Ar-CH)	7.24-8.31 (m,9H,Ar-H),5.47 (s,3H,NH),2.67(s,1H,CH), 3.7 (s,6H,N-Me ₂)
3h	Ph	p-NO ₂	164	71	C ₁₇ H ₁₃ N ₅ S ₂ O ₂	53.26	3.39	18.27	3382 (NH), 765 (C-S), 3062 (Ar-CH), 1435 (C-NO ₂)	7.27-8.35 (m,9H,Ar-H),5.54 (s,3H,NH),2.72 (s,1H,CH)

Antifungal Activity: The compounds (3a-h) were screened for their antifungal activity against *A. niger* and *A. flavus* by known method¹³ at the three concentrations viz. 1000, 100 and 10 ppm. The screening data of compounds are listed in **Table 2**. Results were compared with commercial fungicide miconazole tested under similar conditions.


The percentage inhibition has been calculated by the formula

$$\% \text{ inhibition} = (C-T) \times 100/C$$

Where C and T are a diameter (in mm) of fungus colony in control and treated plates respectively.

TABLE 2: SCREENING DATA OF COMPOUNDS

Comp. no.	R	X	<i>A. niger</i> Concentration (ppm) used			<i>A. flavus</i> Concentration (ppm) used		
			1000	100	10	1000	100	10
3a	H	H	69.7	53.2	35.7	67.2	56.5	32.1
3b	H	p-OMe	67.9	51.8	36.3	68.3	56.8	33.2
3c	H	p-N(Me) ₂	66.5	53.8	35.4	67.1	56.3	32.0
3d	H	p-NO ₂	73.2	57.6	39.0	70.0	57.1	34.8
3e	Ph	H	67.1	56.5	32.0	64.3	50.1	32.8
3f	Ph	p-OMe	67.2	53.5	34.0	65.9	52.3	34.2
3g	Ph	p-N(Me) ₂	65.8	51.4	32.1	64.7	52.1	33.8
3h	Ph	p-NO ₂	70.8	57.3	36.7	72.3	53.4	34.6



RESULT AND DISCUSSIONS: It is observed from the antifungal screening data that most of the compounds have significant toxicity at 1000 ppm, but their toxicity decreases upon dilution **Table 2**. The antifungal activity of the compounds is due to the presence of biolabile groups N=C-S, and N-N=C present in the nuclei of the compound and slight variation in their antifungal activity is due to OCH₃, N-(Me)₂ and NO₂ group attached with benzene nucleus. All the synthesized compounds show antifungal activity against *A. niger* and *A. flavus*.

CONCLUSION: The growth of both fungi *A. niger* and *A. flavus* are inhibited to some extent by all the synthesized compounds. The compounds having a p-nitro group in each set of derivatives are more active. The compounds having p-nitrophenyl and p-methoxy phenyl groups in thiazinopyrazole enhanced the fungicidal activity against both the test fungi *A. niger* and *A. flavus*.

ACKNOWLEDGEMENT: The authors are very grateful to Dr. U. P. Singh (Principal), Dr. Daroga Singh (Ex-Head), Department of Chemistry

How to cite this article:

Chaubey B, Singh VK, Singh D, Verma RK and Yadav VK: Synthesis of some novel 2-substituted imino-7-substituted phenyl-3,7-dihydro-2h-[1, 3]thiazolo[5, 4-e]-thiazin-5-amine as antifungal activity. Int J Life Sci & Rev 2015; 1(1): 20-22. doi: 10.13040/IJPSR.0975-8232.IJLSR.1(1).20-22.

T.D.P.G. College, Jaunpur for providing necessary facilities.

CONFLICT OF INTEREST: Nil

REFERENCES:

- Gupta V and Kant V: A review on biological activity of imidazole and thiazole moieties and their derivatives. Science International 2013; 1: 253-60.
- Narayana B, Raj KKV, Ashalatha BV, Kumari NS and Sarojini BK: Synthesis of some new 5-(2-substituted-1, 3-thiazol-5-yl)-2-hydroxy benzamides and their 2-alkoxy derivatives as possible antifungal agents. European Journal of Medicinal Chemistry 2004; 39: 867-72.
- Mahapatra GN: Non metallic fungicides. Natue 1956; 177: 938.
- Kennedy AD and Summers AL: Chemical constitution and activity of herbicides. Journal of Heterocyclic Chemistry 1981; 18: 409.
- Holla BS, Prasanna CS, Poojary B, Rao KS, Shridhara K and Bhat UG: Synthesis and insecticidal activity of some 1, 3, 4-oxadiazoles derived from 2-chloropyridine-5-acetic acid. Indian Journal of Chemistry 2004; 43(4): 864-8.
- Tiwari N and Nizamuddin: Synthesis of some 1-aryol-3-methyl-4-substituted-phenyl-6-imino-4, 7-dihydro-1, 3-thiazino-5, 4-pyrazoles as pesticides. Journal of Indian Chemical Society 1991; 68: 144-46.
- Desai NC, Parekh BR and Thaker AJ: Preparation of some important medicinal compounds. Thiosemicarbazones, thiadiazolines, 4-thiazolidinones and 5-arylidine derivatives as antibacterial and tuberculostatic agents. Indian Chemical Society 1987; 64(8): 491.
- Geronikaki A, Litina DH, Zablotskaya A and Segal I: Organosilicon-containing thiazole derivatives as potential lipoxigenase inhibitors and anti-inflammatory agents. Bioinorganic Chemistry and Applications 2007: 1-7.
- Kshirsagar A, Turaskar MP, Kulkarni VM, Dhanashire S and Kadam V: Microwave Assisted Synthesis of Potential Anti Infective and Anticonvulsant Thiosemicarbazones. International Journal of Chemical Technology and Research 2009; 1: 696.
- Karali N, Giirsoy A, Kandemirli F, Shvets N, Kaynak FB., Ozbey S, Kovalishym V and Dimoglo A: Synthesis and structure-antituberculosis activity relationship of 1H-indole-2,3-dione derivatives. Bioorganic Medicinal Chemistry 2007; 15: 5888.
- Rawat R, Tripathi R, Katti SB, Pannecouque C and Declereq E: Design, synthesis, and evaluation of 2-aryl-3-heteroaryl-1, 3-thiazolidin-4-ones as anti-HIV agents. Bioorganic Medicinal Chemistry 2007; 15: 1725.
- Gududura V, Hurth E, Dalton JT and Miller DD: Discovery of 2-arylthiazolidine-4-carboxylic acid amides as a new class of cytotoxic agents for prostate cancer. Journal of Medicinal Chemistry 2005; 48: 2584.
- Horsfall JG: Quantitative bioassay of fungicides in the laboratory. Botanical Review 1945; 11(7): 357-97.