



Received on 16 June 2017; received in revised form, 21 July 2017; accepted, 24 July 2017; published 31 July 2017

ANTI-DIABETIC PROPERTIES OF ASIAN EVERYDAY FOOD

Nadia Wajid* and Fatima Ali

The University of Lahore, Lahore, Pakistan.

ABSTRACT: Diabetes mellitus is indicated by abnormal glycemic control. Blurred vision, retinopathy, dry skin, nerve damage, and kidney damage are among the common complications associated with diabetes mellitus. Diet, exercise, medications, and insulin therapy are considered better therapeutic options for diabetes mellitus. Herbal medicines have been a highly valued source of medicine throughout human history. They are widely used today, indicating that herbs are growing parts of modern high-tech medicines and some of the herbal plants and active chemical constituents have a role in the management of diabetes mellitus. In Asia, many herbs are used in everyday food, which has anti-diabetic effects directly or is effective to improve secondary complications like nephropathy, nephropathy, cardiomyopathy, etc. *Allium cepa* (onion), *Allium sativum* (garlic), *Zingiber officinale* (ginger), *Coriandrum sativum* (coriander) are the common herbs used in Asian everyday food which have been reported to alleviate diabetic conditions. The present review summarizes the benefits of these common herbs in diabetes mellitus.

Keywords: Diabetes, Herbs, Onion, Garlic, Coriander, Ginger

Correspondence to Author:

Nadia Wajid

PhD, IMBB, The University of Lahore, Defense Road Campus Lahore, Pakistan.

E-mail: Nadia.wajid@imbb.uol.edu.pk

INTRODUCTION: Diabetes mellitus is caused by a deficiency of insulin hormone, imperfect insulin act, or both, usually a combination of hyperglycemia and the heterogeneous disorders with high blood glucose level¹. Diabetes mellitus and its management have been recognized since the Middle Ages². The World Health Organization (WHO) declared that around 80% world's population rely on herbal therapies due to lack of side effects³. Many plants produce useful compounds for the production of fungicides, insecticides, and industrial raw materials.

In 1980, (WHO) the World Health Organization recommended researchers to find out the therapeutic ability of multiple herbs⁴. It has been estimated that up to one-third of patients with diabetes mellitus use some form of complementary and alternative medicine. To time, approximately 400 ancient plants have been used to treat diabetes, among which only a few have been found to have medicinal importance for reducing blood glucose in non-insulin dependent diabetic patients⁵. Hypoglycemic activity of different plants' products had been used in traditional medicines all over in the world⁶. Asian food contains some of the herbs which are rich in compounds active for therapy of diabetes. Some of the important food plants have been discussed in the present review.

Onion: Scientific Name: *Allium cepa*
Allium cepa, a spice plant, is commonly known as onion and belongs to the family Liliaceae. Since

<p>QUICK RESPONSE CODE</p> 	<p>DOI: 10.13040/IJPSR.0975-8232.IJLSR.3(7).76-78</p> <p>The article can be accessed online on www.ijlsr.com</p>
<p>DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.IJLSR.3(7).76-78</p>	

ancient times, it has been used traditionally for the treatment of different diseases Akash *et al.*⁷ In agriculture, onion is very commonly cultivated in many countries globally. Onion has multi-dimensional effects on diabetes from lowering of blood glucose, lipids, oxidative stress, and chances of thrombosis^{8,9}.

Allium cepa contains sulfur compounds like S-methyl cysteine and flavonoids (quercetin) which are mainly responsible for the hypoglycemic, hypolipidemic, and anti-oxidative stress activities. Extracts of onion normalize the activities of liver hexokinase, glucose 6-phosphatase, and HMG coenzyme-A reductase⁷. Different components of onion-like Cycloalliin, S-methyl-L-cysteine, S-propyl-L-cysteine sulfoxide, dimethyl trisulfide, and S-methyl-L-cysteine sulfoxide inhibit the formation of oil drop in cells¹⁰.

Quercetin has an antiapoptotic effect for liver cells so effective for diabetic hepatopathy¹¹. Onion has also been shown antithrombotic effect by inhibiting collagen- or arachidonic acid (AA) induced platelet aggregation and collagen-induced AA release response¹².

Garlic:

Scientific Name: *Allium sativum*

Garlic (*Allium Sativum*) has been used for various ailments in herbal medicine. Garlic has controversial researches for therapy of diabetes because of inconsistent data from animal models and improper scientific data from humans. Garlic prevents diabetes progression through its antioxidative, anti-inflammatory, and antiglycative properties¹³.

Garlic extracts have been seriously considering red as medicine in recent years as it can reduce serum lipids, blood pressure, and plasma viscosity¹⁴. Extracts of garlic were found to have anti-diabetic activities as they decrease the level of glucose, serum triglycerides, uric acid, urea, creatinine, complete cholesterol, AST and ALT^{15,16}.

Ginger:

Scientific Name: *Zingiber officinale*

Ginger (*Zingiber officinale*) is considered as an effective therapy against diabetes mellitus as it prevents secondary complications. It improves insulin sensitivity, lipid profile, and reduces CRP

and PGE₂^{17, 18, 19}. It inhibits multiple transcriptional pathways, lipid peroxidation, carbohydrate-metabolizing enzymes, and HMG-CoA reductase and activates antioxidant enzyme activity and low-density lipoprotein (LDL) receptors. So it not only improves insulin sensitivity or synthesis but also protects β -cells of pancreatic islets, reduces fat accumulation, decreases oxidative stress, and increases glucose uptake by the tissues.

It also acts as an antioxidant hence improves nephropathy and diabetic cataract²⁰. Ginger has also been reported to reverse diabetic proteinuria in diabetic rats²¹. Oral supplementation of ginger improves glycemic indices, TAC, and PON-1 activity²² reduces FBS (fasting blood sugar) and HbA1c and improves QUICKI (quantitative insulin sensitivity check index)²³.

Coriander:

Scientific Name: *Coriandrum sativum*

Coriandrum sativum (CS) is a daily used food ingredient in Asia, and it is claimed that CS can be used for many ailments. All parts are used for flavoring or treatment of various disorders. It is rich in lipids (petroselinic acid), essential oil (high in linalool), and many bioactive. It has anti-microbial, anti-oxidant, anti-diabetic, anxiolytic, anti-epileptic, anti-depressant, anti-mutagenic, anti-inflammatory, anti-dyslipidemic, anti-hypertensive, neuroprotective and diuretic properties²⁴.

CS extract has many bioactive compounds like phenolics, flavonoids, steroids, and tannins. It lowers blood glucose, total serum cholesterol, triglycerides, and low-density lipoprotein inhibit lipid peroxidation and scavenges 2, 2-diphenyl-2-picrylhydrazyl free radical. It also increases the activity of antioxidant enzymes such as catalase, superoxide dismutase, and glutathione²⁵. Aqueous extract of CS inhibits gastrointestinal α -glucosidase hence has anti-diabetic properties²⁶. It decreases atherosclerosis, increases cardioprotective indices, and several components of diabetes as metabolic syndrome²⁷.

CONCLUSION: From the above data, it may be concluded that Asian everyday herbs are beneficial against diabetes and its complications. Further studies are required to evaluate the improvement of the efficacy of these herbs.

ACKNOWLEDGEMENT: Authors are thankful to The University of Lahore for providing the funding for conducting studies.

CONFLICT OF INTEREST: Authors declare no conflict of interest.

REFERENCES:

- Pari L and Saravanan R: Antidiabetic effect of diasulin, an herbal drug, on blood glucose, plasma insulin and hepatic enzymes of glucose metabolism in hyperglycaemic rats. *Diabetes, Obesity and Metabolism* 2004; 6: 286-292.
- Patlak M: New weapons to combat an ancient disease: Treating diabetes. *Federation of American Society for Experimental Biology* 2002; 16(14): 1853-1857.
- Geoffrey CK: Medicinal plants and control of parasites. *Trans R Soc Trop Med Hyg* 1996; 90: 605-609.
- Anonymous: Diabetes mellitus. *Diabet Med* 1995; 4: 15-19.
- Bailey CJ and Day C: Traditional plant medicines as treatments for diabetes. *Diabetes Care* 1989; 12: 553-564.
- Yaniv Z, Dafni A, Friedman J and Palevitch D: Plants used for the treatment of diabetes in Israel. *Journal Ethnopharmacol* 1987; 19: 145-151.
- Akash MS, Rehman K and Chen S: Spice plant *Allium cepa*: dietary supplement for treatment of type 2 diabetes mellitus. *Nutrition* 2014; 30(10): 1128-37.
- Mathew PT and Augusti KT: Hypoglycemic effects of onion, *Allium cepa* Linn., on diabetes mellitus- a preliminary report. *Indian Journal of Physiology and Pharmacology* 1975; 19: 213-217.
- Bang MA, Kim HA and Cho YJ: Alterations in the blood glucose, serum lipids and renal oxidative stress in diabetic rats by supplementation of onion (*Allium cepa* Linn). *Nutr Res Pract* 2009; 3(3): 242-6.
- Yoshinari O, Shiojima Y and Igarashi K: Anti-obesity effects of onion extract in Zucker diabetic fatty rats. *Nutrients* 2012; 4(10): 1518-26.
- Bakhshaeshi M, Khaki A, Fathiazad F, Khaki AA and Ghadamkheir E: Anti-oxidative role of quercetin derived from *Allium cepa* on aldehyde oxidase (OX-LDL) and hepatocytes apoptosis in streptozotocin-induced diabetic rat. *Asian Pac J Trop Biomed* 2012; 2(7): 528-31.
- Jang EK, Azzam JE, Dickinson NT, Davidson MM and Haslam RJ: Roles for both cyclic GMP and cyclic AMP in the inhibition of collagen-induced platelet aggregation by nitroprusside. *Brit J Haematol* 2002; 117: 664-675.
- Liu CT, Sheen LY and Lii CK: Does garlic have a role as an anti-diabetic agent? *Mol Nutr Food Res* 2007; 51(11): 1353-64.
- Ashraf R, Aamir K, Shaikh AR and Ahmed T: Effects of garlic on dyslipidemia in patients with type 2 diabetes mellitus. *J Ayu Med Coll Abbottabad* 2005; 17(3): 60-64.
- Kumari K, Mathew BC and Augusti KT: Anti-diabetic and hypolipidemic effects of S-methyl cysteine sulfoxide isolated from *Allium cepa* Linn. *Indian J Biochem Biophys* 1995; 32(1):49-54.
- Kumari K and Augusti KT: Lthe lipid-lowering effect of S-methyl cysteine sulfoxide from *Allium cepa* Linn in high cholesterol diet fed rats. *J Ethnopharmacol* 2007; 109(3): 367-71.
- Akhani SP, Vishwakarma SL and Goyal RK: Anti-diabetic activity of *Zingiber officinale* in streptozotocin-induced type I diabetic rat. *J Pharm Pharmacol* 2004; 56(1): 101-5.
- Mahluji S, Attari VE, Mobasseri M, Payahoo L, Ostadrahimi A and Golzari SE: Effects of ginger (*Zingiber officinale*) on plasma glucose level, HbA1c and insulin sensitivity in type 2 diabetic patients. *Int J Food Sci Nutr* 2013; 64(6): 682-6.
- Arablou T, Aryaeian N, Valizadeh M, Sharifi F, Hosseini A and Djalali M: The effect of ginger consumption on glycemic status, lipid profile and some inflammatory markers in patients with type 2 diabetes mellitus. *Int J Food Sci Nutr* 2014; 65(4): 515-20.
- Akash MS, Rehman K, Tariq M, Chen S: *Zingiber officinale* and Type 2 diabetes mellitus: Evidence from experimental studies. *Crit Rev Eukaryot Gene Expr* 2015; 25(2): 91-112.
- Al-Amin ZM, Thomson M, Al-Qattan KK, Peltonen-Shalaby R and Ali M: Anti-diabetic and hypolipidaemic properties of ginger (*Zingiber officinale*) in streptozotocin-induced diabetic rats. *Br J Nutr* 2006; 96(4): 660-6.
- Shidfar F, Rajab A, Rahideh T, Khandouzi N, Hosseini S, and Shidfar S: The effect of ginger (*Zingiber officinale*) on glycemic markers in patients with type 2 diabetes. *J Complement Integr Med* 2015; 12(2): 165-70.
- Mozaffari-Khosravi H, Talaei B, Jalali BA, Najarzadeh A and Mozayan MR: The effect of ginger powder supplementation on insulin resistance and glycemic indices in patients with type 2 diabetes: a randomized, double-blind, placebo-controlled trial. *Com Ther Med* 2014; 22(1): 9-16.
- Sahib NG, Anwar F, Gilani AH, Hamid AA, Saari N and Alkharfy KM: Coriander (*Coriandrum sativum* L.): a potential source of high-value components for functional foods and nutraceuticals-a review. *Phytother Res* 2013; 27(10): 1439-56.
- Sreelatha S and Inbavalli R: Antioxidant, anti-hyperglycemic, and anti-hyperlipidemic effects of *Coriandrum sativum* leaf and stem in alloxan-induced diabetic rats. *J Food Sci* 2012; 77(7): T119-23.
- Brindis F, González-Andrade M, González-Trujano ME, Estrada-Soto S and Villalobos-Molina R: Postprandial glycaemia and inhibition of α -glucosidase activity by aqueous extract from *Coriandrum sativum*. *Nat Prod Res* 2014; 28(22): 2021-5.
- Aissaoui A, Zizi S, Israili ZH and Lyoussi B: Hypoglycemic and hypolipidemic effects of *Coriandrum sativum* L. in Merionesshawi rats. *J Ethnopharmacol* 2011; 137(1): 652-61.

How to cite this article:

Wajid N and Ali F: Anti-diabetic properties of Asian everyday food. *Int J Life Sci & Rev* 2017; 3(7): 76-78. doi: 10.13040/IJPSR.0975-8232.IJLSR.3(7).76-78.

All © 2015 are reserved by International Journal of Life Sciences and Review. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)